

**Appendix 1** to Otti A, Guendel H, Henningsen P, et al. Functional network connectivity of pain-related resting state networks in somatoform pain disorder: an exploratory fMRI study. *J Psychiatry Neurosci* 2012.

DOI: 10.1503/jpn.110187

Copyright © 2012 The Author(s) or their employer(s). To receive this resource in an accessible format, please contact us at [cmajgroup@cmaj.ca](mailto:cmajgroup@cmaj.ca).

**Table S1: Brief Pain Inventory item scores for each patient\***

Patient	BPI item													
	1	2	3	4	5	6	8	9A	9B	9C	9D	9E	9F	9G
1	Yes	Yes	5	5	5	2	3	4	5	4	3	7	6	8
2	Yes	Yes	8	3	6	5	8	8	7	7	7	4	4	7
3	Yes	Yes	8	7	7	8	7	8	9	6	7	7	10	9
4	Yes	Yes	10	9	9	10	9	10	4	9	10	9	9	3
5	Yes	Yes	9	4	5	6	5	7	7	0	7	3	1	4
6	Yes	Yes	2	1	2	1	—	2	1	3	3	1	0	1
7	Yes	Yes	4	0	2	1	—	3	1	2	3	1	2	1
8	Yes	Yes	—	—	—	—	—	—	—	—	—	—	—	—
9	Yes	Yes	6	3	5	4	3	5	6	1	8	5	5	7
10	Yes	Yes	10	7	9	9	0	9	9	10	1	5	9	7
11	Yes	Yes	10	7	8	10	0	9	0	4	2	0	2	0
12	Yes	Yes	7	5	6	8	5	4	2	1	4	0	6	1
13	Yes	Yes	5	4	4	5	5	4	3	5	7	4	5	3
14	Yes	Yes	8	6	8	8	7	8	9	8	8	6	8	8
15	Yes	Yes	—	—	—	—	—	—	—	—	—	—	—	—
16	Yes	Yes	8	6	7	7	—	4	2	3	4	2	5	3
17	Yes	Yes	5	1	3	3	9	3	5	3	6	5	5	6
18	Yes	Yes	7	4	6	5	3	6	5	7	3	4	7	5
19	Yes	Yes	5	0	4	3	5	2	3	2	4	1	0	3
20	Yes	Yes	7	3	4	3	5	5	7	0	7	5	3	6
21	Yes	Yes	9	5	7	7	2	8	7	7	8	7	6	7

BPI = Brief Pain Inventory (Radbruch L, Loick G, Kiencke P, et al. Validation of the German version of the Brief Pain Inventory. *J Pain Symptom Manage* 1999;18:180-7.)

\*Missing data are indicated with an em-dash. BPI items are as follows: 1 = pain within the last week, 2 = pain today, 3 = pain at its worst during the last week, 4 = pain at its least during the last week, 5 = pain on the average, 6 = pain right now, 8 = pain relief by therapy, 9A = impairment of general activity, 9B = impairment of mood, 9C = impairment of walking ability, 9D = impairment of normal work, 9E = impairment of relations with other people, 9F = impairment of sleep, 9G = impairment of enjoyment of life.