

Amygdala and insula response to emotional images in patients with generalized social anxiety disorder

Sabin G. Shah, BS; Heide Klumpp, PhD; Mike Angstadt, BS; Pradeep J. Nathan, PhD;
K. Luan Phan, MD

Shah, Klumpp, Angstadt, Phan — Department of Psychiatry, University of Michigan, Ann Arbor, Mich.; Nathan — Department of Psychiatry, University of Cambridge, UK; Phan — Mental Health Service, VA Ann Arbor Healthcare System, Ann Arbor, Mich.

Background: Functional brain imaging studies have demonstrated amygdala and insula hyper-reactivity to probes of social threat in participants with generalized social anxiety disorder (gSAD). The amygdala and insula are known to serve broad functions in emotional processing, including integration of affective information. However, few studies have examined brain responses in socially anxious participants during general emotional processing. We examined brain response to emotionally evocative images in patients with gSAD and matched healthy controls. **Methods:** Eleven patients with gSAD who were not taking psychotropic medications and did not have psychiatric comorbidities and 11 matched healthy controls underwent functional magnetic resonance imaging while viewing blocks of emotionally salient (positive, negative, neutral) pictures. **Results:** Participants with gSAD exhibited enhanced bilateral amygdala and insula reactivity to negative (v. neutral) images compared with healthy controls who did not exhibit enhanced reactivity. Within the gSAD group, the extent of amygdala activation was correlated with social anxiety severity, whereas the extent of insula activation was correlated with trait anxiety. **Limitations:** The small sample size may have limited our ability to detect group differences in other relevant brain regions and in behavioural measures. **Conclusion:** In addition to prior findings of probes of social information processing, our findings suggest that the amygdala and insula responses are hyper-reactive to general emotional images with negative emotional content and that these brain regions may play divergent roles in their representation of different phenotypes.

Introduction

Generalized social anxiety disorder (gSAD), also known as social phobia, is characterized by an exaggerated fear of negative scrutiny across interpersonal interactions. This exaggerated fear response may be in part due to excessive attention to negative stimuli^{1,2} and/or deficits in processing positive^{3,4} or ambiguous^{5,6} interpersonal stimuli. The neural correlates of this fear response have been extensively investigated by social probes of potential threat (e.g., harsh faces, public speaking, verbal criticism),⁷⁻¹² and a recent meta-analysis of functional neuroimaging studies¹³ confirmed that heightened activation of the amygdala and insula is a robust and consistent finding in social phobia. The amygdala and insula are known to serve broad functions in emotional processing, including mind-body integration of affective information,^{14,15} and their increased sensitivity to negative social stimuli has been specifically implicated as a neural marker of social anxiety severity.^{8,11,12,16}

To date, we are not aware of any studies that have examined brain responses in socially anxious participants during general (nonsocial, e.g., not faces or public speaking) emotional processing. We compared brain response to complex and emotionally evocative images in patients with gSAD and matched healthy controls. We employed stimuli from the International Affective Picture System (IAPS), a standardized set of images known to reliably evoke acute and transient changes in emotional experience and arousal as well as central (e.g., amygdala and insula) indices of emotional reactivity.¹⁷ Specifically, we hypothesized that amygdala and insula reactivity in patients with gSAD would be exaggerated compared with healthy controls in response to general emotional probes of negative, but not positive, affect. Although it may be argued that anxious temperament and nonspecific/general anxiety have overlapping characteristics, the Spielberger Trait Anxiety Inventory (STAI-trait)¹⁸ measures anxiety proneness as a stable temperamental/personality trait

Correspondence to: Dr. K.L. Phan, Department of Psychiatry, University of Michigan, Rachel Upjohn Bldg., Room 2751, Box 6765, 4250 Plymouth Rd., Ann Arbor, MI 48109-2700; fax 734 936-7868; luan@umich.edu

J Psychiatry Neurosci 2009;34(4):296-302.

Submitted Jul. 29, 2008; Revised Oct. 24, Dec. 18, 2008; Accepted Dec. 23, 2008

(e.g., individual differences in the likelihood that people would experience state anxiety in a stressful situation in their lifetimes), whereas the Hamilton Anxiety Rating Scale (HAM-A)¹⁹ and Liebowitz Social Anxiety Scale (LSAS)²⁰ measure state transitory levels of current generalized, non-specific anxiety (e.g., anxiety symptoms over the past week) and social anxiety, and thus represent divergent constructs.²¹ Interestingly, in individuals with social anxiety or with high anxiety proneness, the extent of their amygdala and/or insula reactivity is positively associated with their levels of trait anxiety and social anxiety but not with their levels of nonspecific anxiety.^{8,11,22} Thus, we further hypothesized that the magnitude of amygdala and/or insula reactivity would be associated with the level of symptom severity and/or anxious temperament, but not with nonspecific/general anxiety.

Methods

Participants

We recruited participants with gSAD based on DSM-IV criteria, as confirmed with the Structured Clinical Interview for DSM-IV (SCID-IV)²³ with additional probes from the clinician-administered LSAS,²⁰ from the community by advertisement. In addition to the LSAS, participants with gSAD also completed questionnaires to measure trait anxiety and clinical symptomatology measures, including the Spielberger State-Trait Anxiety Inventory (STAI),¹⁸ Beck Depression Inventory (BDI)²⁴ and the Hamilton Rating Scales for Depression (HAM-D)²⁵ and Anxiety (HAM-A).¹⁹ Severity of social anxiety, trait anxiety and general/nonspecific anxiety were represented by scores on LSAS, STAI-trait and HAM-A, respectively. We recruited matched healthy controls free of prior or current medical, neurologic and psychiatric illness, as confirmed by SCID-IV and medical evaluation by a physician (K.L.P.). Trained clinicians and a board-certified psychiatrist (K.L.P.) performed clinical assessments using SCID-based DSM-IV criteria and clinical judgment based on direct patient interviews and review of all symptom ratings scores. All participants provided written informed consent, and the local university institutional review board approved our study.

Experimental task

We presented 60 IAPS images showing any of 3 affect conditions (i.e., negative, positive, neutral) (Box 1) in 20-second blocks (5 images/block, 2 blocks of each valence/run), alternating with 20-second baseline blocks of blank, greyscale images (6 blocks/run), over 2 functional runs of 4 minutes each. We did not repeat images, and we counterbalanced block order. We selected the negative (i.e., unpleasant/aversive) and positive (i.e., pleasant) images based on normative ratings of low and high valence (pleasantness), respectively, with similar levels of subjective arousal.

Before scanning, we introduced participants to the task and performed a short practice set containing distinct IAPS images that we did not later present in the scanner. During

functional magnetic resonance imaging (fMRI) scanning, we instructed participants to identify the general valence of each image (i.e., negative, positive or neutral) by button-press as soon as they were able to judge the image. After scanning, participants rated each image on a 9-point scale for valence and arousal (1 = most unpleasant/least arousing; 9 = most pleasant/most arousing).

MRI acquisition and analysis

We performed all scanning with blood oxygen level-dependent (BOLD) whole-brain fMRI on a 3.0 T GE Signa

Box 1: International Affective Picture System (IAPS) images* by category and number

	Category, no.		
	Neutral	Positive	Negative
	1121	1440	1300
	1313	1463	2053
	1390	1710	2205
	1670	1920	2692
	1945	1999	2700
	2214	2058	2710
	2280	2070	2730
	2372	2160	2800
	2381	2165	2900
	2383	2340	3030
	2440	2550	3060
	2480	2660	3150
	2485	4220	3180
	2487	4250	3220
	2514	4572	3266
	2570	4599	3300
	2575	4608	6020
	2580	4660	6230
	2749	5450	6312
	2840	5470	6313
	2850	5480	6560
	2870	5831	6570
	2880	5982	7380
	4605	7282	8230
	5395	7330	9000
	5531	7352	9006
	5535	7502	9040
	5740	8034	9050
	6150	8080	9220
	7002	8120	9290
	7010	8161	9300
	7130	8185	9410
	7185	8300	9440
	7217	8370	9470
	7233	8380	9560
	7500	8496	9561
	7595	8500	9570
	9070	8501	9611
	9210	8531	9800
	9700	8540	9910

*The mean (and standard deviation [SD]) ratings for valence were 5.0975 (0.464459) for neutral, 7.4415 (0.504658) for positive and 2.4025 (0.544241) for negative images and the mean (and SD) ratings for arousal were 3.3985 (0.72151) for neutral, 5.4475 (0.820193) for positive and 5.77225 (0.881207) for negative images.

System (General Electric) using a standard radiofrequency coil and associated software (LX 8.3, Neuro-optimized gradients; General Electric). We acquired whole-brain functional scans using a T_2 -weighted reverse spiral gradient-recall echo sequence (echo time [TE] 25 ms, repetition time [TR] 2000 ms, 64×64 matrix, flip angle 77° , field of view [FOV] 24 cm, 3.75 mm^2 inplane voxels, 30 contiguous 5-mm axial slices per volume) optimized to minimize susceptibility artifacts in the regions of interest (ROIs).²⁶ We also acquired a high-resolution T_1 -weighted scan (3-dimensional magnetization prepared rapid gradient echo, or 3D-MPRAGE; TR 25 ms; min TE; FOV 24 cm; slice thickness 1.5 mm) for anatomic localization.

Functional MRI data preprocessing

Data from all participants met the criteria for quality with minimal motion correction (< 3 mm displacement in any one direction); we included these data in our analyses. We discarded the first 4 volumes from each run to allow for T_1 equilibration effects. We preprocessed and analyzed the data using statistical parametric mapping (SPM5; Wellcome Department of Cognitive Neurology). We spatially realigned the scans to the first scan in each run to correct for head motion, warped (nonlinear) to an echo-planar image (EPI) template in Montreal Neurologic Institute (MNI) space, resampled them to 2-mm^3 voxels and smoothed them with an 8-mm^3 kernel to maximize signal and minimize residual differences in neuroanatomy. We combined the general linear model applied to the time series with the canonical hemodynamic response function and with a 128-second high-pass filter.²⁷

Statistical analysis

We performed conventional data processing analyses based on the general linear model using SPM5 software (Wellcome Department of Cognitive Neurology). We generated linear contrasts of interest (negative v. neutral, positive v. neutral) for each participant, and entered them into a second-level random effects model and 2-sample Student t tests to examine between-group differences in brain activation. Within ROIs defined a priori (i.e., amygdala and insula), we set the significance threshold at $p_{\text{uncorrected}} < 0.005$, with an extent of greater than 10 contiguous voxels to balance between type I and type II errors.²⁸ Outside the ROIs defined a priori, we set the threshold at $p < 0.05$, corrected for multiple comparisons using a false discovery rate method.²⁹ Following analyses, we identified the location of activation foci using MARINA software (www.bion.de/index.php?title=MARINA)³⁰ based on the atlas by Tzourio-Mazoyer and colleagues.³¹ Additionally, we extracted parameter estimates (i.e., β weights, arbitrary units), an index of activation signal change, from 10-mm diameter spherical volumes around the peak voxels within bilateral insula and the amygdala using MarsBaR (<http://marsbar.sourceforge.net>), and we used 2-tailed Student t tests for between-group comparisons and calculated Pearson correlation coefficients with symptom/trait measures (i.e., LSAS, STAI, HAM-A); we set significance at $p < 0.05$, 2-tailed.

Results

Participants

We enrolled 22 right-handed participants (11 with gSAD and 11 controls) in our study. The groups did not differ significantly in age (mean 27.45, standard deviation [SD] 8.96, range 19–49 yr in the gSAD group v. mean 30.55, SD 7.69, range 21–49 yr in the control group; $p = 0.40$) or sex (8 men and 3 women in the gSAD group v. 6 men and 5 women in the control group; $p = 0.38$). We were able to collect data from the STAI, BDI, HAM-D and HAM-A questionnaires for all participants with gSAD and 7 of the 11 controls. Mean (SD) self-report scores on LSAS, STAI, HAM-A, HAM-D and BDI and between-group comparisons are reported in Table 1. Of note, none of these 3 scores correlated with one another within the gSAD or control groups (Pearson correlations, all $p > 0.05$).

All participants were free of psychoactive medications at the time of scanning and urine toxicology screens at the time of scanning were negative for all participants. None had current or recent (previous 6 mo) depressive episodes, substance abuse or a history of autism/pervasive developmental disorders, mental retardation or significant medical or neurologic illness, as verified by the SCID-IV and a medical history screening. Two patients with gSAD had concurrent generalized anxiety disorder; however, it did not precede the onset of social anxiety symptoms, nor was it more clinically salient than gSAD.

Behavioural results

Behavioural results are summarized in Table 2. We observed no significant group-by-condition interactions across valence categories; participants with gSAD did not differ from controls on ratings of valence or arousal for negative, positive or neutral picture conditions. Across participants, there was a significant main effect of condition for both valence and arousal ratings. Follow-up Student t tests revealed that, relative to neutral pictures, negative pictures were rated more

Table 1: Self-report measures and between-group comparisons* for 11 participants with generalized social anxiety disorder and 11 healthy controls

Measure	Group; mean (SD)		t_{16} value	Cohen d
	gSAD	Control		
LSAS	75.91 (14.87)	13.71 (14.26)	8.79†	4.27
STAI-state	41.73 (11.21)	30.14 (7.49)	2.40‡	1.22
STAI-trait	46.09 (10.44)	30.42 (4.72)	3.71§	1.94
BDI	9.45 (5.72)	2.29 (3.30)	3.00§	1.53
HAM-D	4.45 (5.94)	0.86 (1.57)	1.55	0.83
HAM-A	4.00 (3.10)	0.71 (1.11)	2.67‡	1.41

BDI = Beck Depression Inventory²⁴; gSAD = generalized social anxiety disorder; HAM-A = Hamilton Rating Scale for Anxiety²⁵; HAM-D = Hamilton Rating Scale for Depression²⁵; LSAS = Liebowitz Social Anxiety Scale²⁶; SD = standard deviation; STAI = State-Trait Anxiety Inventory.¹⁸

*We performed group comparisons using the Student t test for independent groups.

† $p < 0.001$.

‡ $p < 0.01$.

§ $p < 0.05$.

unpleasant, and positive pictures were rated more pleasant on valence; negative and positive pictures were rated to be equally arousing but more arousing than neutral pictures (data not shown). There were no significant group differences in accuracy or reaction time in the in-scanner task.

Functional MRI results in amygdala and insula

Compared with controls, participants with gSAD exhibited more differential activation to negative (> neutral) images in the right amygdala, extending into the superior temporal

Table 2: Subjective ratings and group-by-condition analysis of variance for 11 participants with generalized social anxiety disorder and 11 healthy controls

Rating	Group; mean (SD)*		Group	Condition	Interaction
	gSAD	Control	$F_{1,20}$ value	$F_{2,40}$ value	$F_{2,40}$ value
Valence					
Negative	2.45 (0.35)	2.1 (0.64)	2.98	323.15†	0.74
Positive	6.78 (0.82)	6.84 (0.53)			
Neutral	5.25 (0.42)	4.99 (0.39)			
Arousal					
Negative	6.34 (1.12)	6.35 (1.58)	0.13	44.30†	0.30
Positive	5.46 (1.08)	5.49 (1.32)			
Neutral	3.30 (1.30)	2.85 (1.56)			
In-scan accuracy, %					
Negative	82.3 (20.4)	88.6 (14.0)	0.58	3.94‡	1.16
Positive	78.2 (22.6)	73.2 (17.5)			
Neutral	66.4 (16.6)	76.4 (16.7)			
In-scan reaction time, ms					
Negative	1605.97 (351.60)	1519.10 (290.56)	0.99	4.42‡	0.97
Positive	1689.97 (289.06)	1482.00 (255.91)			
Neutral	1749.57 (374.07)	1694.94 (323.63)			

gSAD = generalized social anxiety disorder; SD = standard deviation.

*Unless indicated otherwise.

† $p < 0.001$.

‡ $p < 0.05$.




Fig. 1: Bilateral amygdala hyperactivity in response to negative (> neutral) images in participants with generalized social anxiety disorder (gSAD). (A) Whole-brain, voxel-wise *t*-map image of group differences depicting greater activation in the bilateral amygdala in patients with gSAD compared with controls. (B) Scatterplot of signal change (negative > neutral images) within the left and right amygdala volumes for each participant with gSAD (red) and control (green). The *t*-map is superimposed on coronal sections of a canonical brain image (at *y*-plane coordinates 4 on the right [R], and -10 on the left [L], using the Montreal Neurologic Institute atlas).

gyrus, and the left amygdala, extending into the hippocampus (peak voxel [36,4,-28], $Z = 3.84$, 544 mm^3 in the right v. peak voxel [-26,-10,-14], $Z = 2.86$, 128 mm^3 in the left; Fig. 1). Compared with controls, participants with gSAD also exhibited more differential activation to negative (> neutral) images in bilateral insula, extending into the putamen (peak voxel [36,-6,4], $Z = 2.83$, 136 mm^3 in the right v. peak voxel [48,8,6], $Z = 2.81$, 208 mm^3 also in the right v. peak voxel [-26,10,6], $Z = 2.88$, 208 mm^3 in the left; Fig. 2). The 2 groups did not differ in amygdala nor insula response to positive (> neutral) images. Of note, we observed no group differences in the amygdala or insula in the neutral > blanks contrast. Activation signal (parameter estimates, mean [SD]) was greater in the amygdala among participants with gSAD than controls ($0.75 [0.52]$ v. $-0.06 [0.42]$, $t_{20} = 4.04$, $p = 0.001$, Cohen $d = 1.71$ in the right amygdala and $0.59 [0.57]$ v. $-0.05 [0.51]$, $t_{20} = 2.77$, $p = 0.012$, Cohen $d = 1.18$ in the left amygdala). Activation signal (parameter estimates, mean [SD]) was also greater in the insula among participants with gSAD than controls ($0.45 [0.55]$ v. $-0.13 [0.37]$, $t_{20} = 2.94$, $p = 0.008$, Cohen $d = 1.24$ in the right insula and $0.20 [0.35]$ v. $-0.18 [0.24]$, $t_{20} = 3.00$, $p = 0.007$, Cohen $d = 1.27$ in the left insula). Among participants with gSAD, the magnitude of BOLD response to negative (> neutral) images in the right amygdala was associated with the intensity of social anxiety symptoms (LSAS_{total}; $r_9 = 0.62$, $p = 0.042$) but not with any other symptom/trait measures; response in the left amygdala was not correlated with any symptom measures. There were no significant correlations between insula activation and any symptom measures at the $p < 0.05$ level. However, the magnitude of both right and left

insula activation was correlated with trait anxiety measures at the trend level ($r_9 = 0.55$, $p = 0.08$ in the right insula and $r_9 = 0.55$, $p = 0.08$ in the left insula).

Functional MRI results outside a priori ROIs

Outside of the insula and amygdala, there were no ROIs that survived correction for multiple comparisons for gSAD > control or control > gSAD in either negative or positive (v. neutral) pictures.

Discussion

To our knowledge, ours is the first study to evaluate the neural correlates of general emotional processing in patients with gSAD. The present study provided evidence of greater differential amygdala and insula reactivity in participants with gSAD, relative to matched controls, in response to images with negative, but not positive, emotional content. Moreover, we observed that the magnitude of this reactivity in the amygdala was associated with severity of social anxiety, whereas insula reactivity was associated at a trend level with trait anxiety. These findings add to the growing literature implicating the amygdala and insula as key components to a common neural marker of SAD.

Exaggerated amygdala and insular reactivity to negative emotional stimuli has also been observed in a number of prior functional brain imaging studies of harsh (e.g., emotionally negative) faces,^{7,8,11,12} unpleasant anticipation before public speaking^{9,32} and verbal criticism.¹⁰ Similar to our




Fig. 2: Bilateral insula hyperactivity in response to negative (> neutral) images in participants with generalized social anxiety disorder (gSAD). (A) Whole-brain, voxel-wise t -map image of group differences depicting greater activation in bilateral insula in participants with gSAD compared with controls. (B) Scatterplot of signal change (negative > neutral images) within the left and right insula volumes for each participant with gSAD (red) and control (green). The t -map is superimposed on axial sections of a canonical brain image (at z -plane coordinates 6 on the right [R], and 8 on the left [L], using the Montreal Neurologic Institute atlas).

results, differences between gSAD and comparison participants in amygdala and insula response to positively valenced stimuli (e.g., happy faces, verbal praise) have not been observed^{7,8,10,11} in accord with behavioural models linking social anxiety and negative (but not positive) bias in socio-emotional information processing.³³ It should be noted that Straube and colleagues³ have observed amygdala hyper-reactivity in participants with SAD to both angry and happy faces. Our findings extend evidence of enhanced amygdala and insula sensitivity to general emotional images with general negative content (e.g., unpleasant/aversive pictures). The amygdala and insula are heavily interconnected³⁴ and are believed to play a concerted role in the regulation of autonomic responses, processing negative affective experiences, and making social judgments (e.g., trustworthiness, approachability) from facial expressions of emotion,^{14,15,35,36} making them highly plausible neural substrates in the pathophysiology of generalized social phobia.^{13,14}

We found that the magnitude of BOLD response to negative emotional images in the amygdala among participants with gSAD was correlated with their levels of social anxiety severity, but not with anxiety temperament (e.g., trait anxiety) or nonspecific (e.g., nonsocial) anxiety symptoms. This is consistent with a number of prior studies^{8,11,12,16} and suggests that amygdala reactivity may be a useful biomarker with clinical application; amygdala hyper-reactivity is normalized by and predicts effective treatment of social anxiety.^{37,38} In addition, insula reactivity was related to levels of trait anxiety at a trend level but not to social anxiety severity. This is consistent with prior findings showing that anxiety-prone (e.g., high-trait anxious) individuals exhibit greater insular reactivity to emotional faces than anxiety-normative controls, and their levels of trait anxiety predict the extent of insula activation.²²

Limitations

Our study has limitations worth discussion. Although similar to prior functional neuroimaging studies of gSAD, our sample size was small, which may have limited our ability to detect group differences in other brain regions or in subjective ratings (e.g., type II error). We were unable to directly test for differences between groups on processes that may have contributed to the observed results (e.g., IQ, habituation to images) because of the study design. This is an important issue that will need to be addressed in future studies. Additionally, our controls did not show amygdala activity in response to negative images relative to neutral images, similar to results of some^{17,39,40} but not all prior studies.⁴¹⁻⁴³ Several factors may contribute to the inconsistency of these findings. Prior evidence of amygdala response to negative images in controls was generated from contrasts with nonimages^{41,42} and involved passive viewing of images⁴¹ or implicit processing of negative images,^{42,43} whereas the task in our study involved explicit cognitive appraisal/labelling of the emotional content, which may have diminished amygdala reactivity to unpleasant pictures in healthy controls.⁴⁴ Moreover, a number of studies, including some conducted in our laboratory,

have shown that the amygdala is activated when viewing neutral pictures^{17,39} and thereby diminishes the activation difference observable in the negative versus neutral contrast.^{17,39}

Despite limitations, our data suggest that the amygdala and insula may be key brain regions in the common final pathway in the neuropathogenesis of SAD, but that they may play divergent roles in their representation of phenotypic markers. It may be that amygdala reactivity to probes of negative emotions or social threat reflects the disease process of social anxiety rather than a vulnerability marker that increases risk for disease, whereas insula reactivity may be better linked to a temperamental risk, or diathesis, toward the development of social or another anxiety disorder. Therefore, our findings support the inclusion of the amygdala and insula as critical components of a brain-based model of SAD.

Acknowledgements: This research was supported by a Seed Grant from the Brain Research Foundation (K.L.P.) and an NIH grant MH076198 (K.L.P.). We thank Rose McCarron for her assistance in participant recruitment and study coordination.

Competing interests: None declared.

Contributors: Dr. Phan designed the study. Mr. Angstadt acquired the data, which all authors analyzed. Mr. Shah and Drs. Klumpp and Phan wrote the article, which Mrs. Angstadt and Shah and Drs. Nathan and Phan reviewed. All authors gave final approval for publication.

References

1. Mogg K, Bradley BP. Selective orienting of attention to masked threat faces in social anxiety. *Behav Res Ther* 2002;40:1403-14.
2. Mogg K, Philippot P, Bradley BP. Selective attention to angry faces in clinical social phobia. *J Abnorm Psychol* 2004;113:160-5.
3. Straube T, Mentzel HJ, Miltner WH. Common and distinct brain activation to threat and safety signals in social phobia. *Neuropsychobiology* 2005;52:163-8.
4. Moser JS, Hajcak G, Huppert JD, et al. Interpretation bias in social anxiety as detected by event-related brain potentials. *Emotion* 2008;8:693-700.
5. Cooney RE, Atlas LY, Joormann J, et al. Amygdala activation in the processing of neutral faces in social anxiety disorder: Is neutral really neutral? *Psychiatry Res* 2006;148:55-9.
6. Yoon KL, Zinbarg RE. Interpreting neutral faces as threatening is a default mode for socially anxious individuals. *J Abnorm Psychol* 2008;117:680-5.
7. Stein MB, Goldin PR, Sareen J, et al. Increased amygdala activation to angry and contemptuous faces in generalized social phobia. *Arch Gen Psychiatry* 2002;59:1027-34.
8. Phan KL, Fitzgerald DA, Nathan PJ, et al. Association between amygdala hyperactivity to harsh faces and severity of social anxiety in generalized social phobia. *Biol Psychiatry* 2006;59:424-9.
9. Tillfors M, Furmark T, Marteinsdottir I, et al. Cerebral blood flow during anticipation of public speaking in social phobia: a PET study. *Biol Psychiatry* 2002;52:1113-9.
10. Blair K, Geraci M, Devido J, et al. The neural response to self-and-other-referential praise and criticism in generalized social phobia (GSP). *Arch Gen Psychiatry* 2008;65:1176-84.
11. Evans KC, Wright CI, Wedig MM, et al. A functional MRI study of amygdala responses to angry schematic faces in social anxiety disorder. *Depress Anxiety* 2008;25:496-505.
12. Goldin PR, Hakimi S, Manber T, et al. Neural bases of social anxiety disorder: emotional reactivity and cognitive regulation during social and physical threat. *Arch Gen Psychiatry*. In press.
13. Etkin A, Wager TD. Functional neuroimaging of anxiety: a meta-analysis of emotional processing in PTSD, social anxiety disorder, and specific phobia. *Am J Psychiatry* 2007;164:1476-88.

14. Paulus MP, Stein MB. An insular view of anxiety. *Biol Psychiatry* 2006;60:383-7.
15. Zald DH. The human amygdala and the emotional evaluation of sensory stimuli. *Brain Res Brain Res Rev* 2003;41:88-123.
16. Killgore WD, Yurgelun-Todd DA. Social anxiety predicts amygdala activation in adolescents viewing fearful faces. *Neuroreport* 2005;16:1671-5.
17. Liberzon I, Phan KL, Decker LR, et al. Extended amygdala and emotional salience: a PET activation study of positive and negative affect. *Neuropsychopharmacology* 2003;28:726-33.
18. Spielberger CD, Gorsuch RL, Lushene R. *Manual for State-Trait Anxiety Inventory*. Palo Alto (CA): Consulting Psychologists Press; 1970.
19. Hamilton M. The assessment of anxiety states by rating. *Br J Med Psychol* 1959;32:50-5.
20. Heimberg RG, Horner KJ, Juster HR, et al. Psychometric properties of the Liebowitz Social Anxiety Scale. *Psychol Med* 1999;29:199-212.
21. Keedwell P, Snaith RP. What do anxiety scales measure? *Acta Psychiatr Scand* 1996;93:177-80.
22. Stein MB, Simmons AN, Feinstein JS, et al. Increased amygdala and insula activation during emotion processing in anxiety-prone subjects. *Am J Psychiatry* 2007;164:318-27.
23. First MB, Spitzer RL, Williams JBW, et al. *Structured Clinical Interview for DSM-IV. Patient Edition (SCID-P)*. Washington (DC): American Psychiatric Press; 1995.
24. Beck AT, Ward CH, Mendelson M, et al. An inventory for measuring depression. *Arch Gen Psychiatry* 1961;4:561-71.
25. Hamilton M. A rating scale for depression. *J Neurol Neurosurg Psychiatry* 1960;23:56-62.
26. Stenger VA, Boada FE, Noll DC. Three-dimensional tailored RF pulses for the reduction of susceptibility artifacts in T*(*)2-weighted functional MRI. *Magn Reson Med* 2000;44:525-31.
27. Friston KJ, Holmes AP, Worsley KJ, et al. Statistical parametric maps in functional imaging: a general linear approach. *Hum Brain Mapp* 1995;2:189-210.
28. Forman SD, Cohen JD, Fitzgerald M, et al. Improved assessment of significant activation in functional magnetic resonance imaging (fMRI): use of a cluster-size threshold. *Magn Reson Med* 1995;33:636-47.
29. Genovese CR, Lazar NA, Nichols T. Thresholding of statistical maps in functional neuroimaging using the false discovery rate. *Neuroimage* 2002;15:870-8.
30. Walter B, Blecker C, Kirsch P, et al. MARINA: an easy to use tool for the creation of masks for region of interest analyses [abstract]. Presented at the 9th International Conference on Functional Mapping of the Human Brain; 2003 Jun. 19-22; New York (NY). Available on CD-Rom in *NeuroImage* 2003;19(2).
31. Tzourio-Mazoyer N, Landeau B, Papathanassiou D, et al. Automated anatomical labeling of activations in SPM using a macroscopic anatomical parcellation of the MNI MRI single-subject brain. *Neuroimage* 2002;15:273-89.
32. Lorberbaum JP, Kose S, Johnson MR, et al. Neural correlates of speech anticipatory anxiety in generalized social phobia. *Neuroreport* 2004;15:2701-5.
33. Bogels SM, Mansell W. Attention processes in the maintenance and treatment of social phobia: hypervigilance, avoidance and self-focused attention. *Clin Psychol Rev* 2004;24:827-56.
34. Augustine JR. Circuitry and functional aspects of the insular lobe in primates including humans. *Brain Res Brain Res Rev* 1996;22:229-44.
35. Craig AD. How do you feel? Interoception: the sense of the physiological condition of the body. *Nat Rev Neurosci* 2002;3:655-66.
36. Winston JS, Strange BA, O'Doherty J, et al. Automatic and intentional brain responses during evaluation of trustworthiness of faces. *Nat Neurosci* 2002;5:277-83.
37. Furmark T, Tillfors M, Marteinsdottir I, et al. Common changes in cerebral blood flow in patients with social phobia treated with citalopram or cognitive-behavioral therapy. *Arch Gen Psychiatry* 2002;59:425-33.
38. McClure EB, Adler A, Monk CS, et al. fMRI predictors of treatment outcome in pediatric anxiety disorders. *Psychopharmacology (Berl)* 2007;191:97-105.
39. Phan KL, Fitzgerald DA, Gao K, et al. Real-time fMRI of cortico-limbic brain activity during emotional processing. *Neuroreport* 2004;15:527-32.
40. Lang PJ, Bradley MM, Fitzsimmons JR, et al. Emotional arousal and activation of the visual cortex: an fMRI analysis. *Psychophysiology* 1998;35:199-210.
41. Britton JC, Taylor SF, Sudheimer KD, et al. Facial expressions and complex IAPS pictures: common and differential networks. *Neuroimage* 2006;31:906-19.
42. Hariri AR, Tessitore A, Mattay VS, et al. The amygdala response to emotional stimuli: a comparison of faces and scenes. *Neuroimage* 2002;17:317-23.
43. Meseguer V, Romero MJ, Barros-Loscertales A, et al. Mapping the appetitive and aversive systems with emotional pictures using a block-design fMRI procedure. *Psicothema* 2007;19:483-8.
44. Taylor SF, Phan KL, Decker LR, et al. Subjective rating of emotionally salient stimuli modulates neural activity. *Neuroimage* 2003;18:650-9.

drcareers
 CMA's career centre for physicians
 Sign up for personalized job alerts.

cma.ca/drcareers

ASSOCIATION MÉDICALE CANADIENNE | CANADIAN MEDICAL ASSOCIATION