

43. Fischl B, Salat DH, van der Kouwe AJ, et al. Sequence-independent segmentation of magnetic resonance images. *Neuroimage* 2004; 23:S69-84.
44. Fischl B, Sereno MI, Dale AM. Cortical surface-based analysis: II: Inflation, flattening, and a surface-based coordinate system. *Neuroimage* 1999;9:195-207.
45. Fischl B, Sereno MI, Tootell RB, et al. High-resolution intersubject averaging and a coordinate system for the cortical surface. *Hum Brain Mapp* 1999;8:272-84.
46. Fischl B, van der Kouwe A, Destrieux C, et al. Automatically parcellating the human cerebral cortex. *Cereb Cortex* 2004;14:11-22.
47. Han X, Jovicich J, Salat D, et al. Reliability of MRI-derived measurements of human cerebral cortical thickness: the effects of field strength, scanner upgrade and manufacturer. *Neuroimage* 2006;32: 180-94.
48. Jovicich J, Czanner S, Greve D, et al. Reliability in multi-site structural MRI studies: effects of gradient non-linearity correction on phantom and human data. *Neuroimage* 2006;30:436-43.
49. Ségonne F, Pacheco J, Fischl B. Geometrically accurate topology-correction of cortical surfaces using nonseparating loops. *IEEE Trans Med Imaging* 2007;26:518-29.
50. Reuter M, Rosas HD, Fischl B. Highly accurate inverse consistent registration: a robust approach. *Neuroimage* 2010;53:1181-96.
51. Ségonne F, Dale A, Busa E, et al. A hybrid approach to the skull stripping problem in MRI. *Neuroimage* 2004;22:1060-75.
52. van Erp TG, Hibar D, Rasmussen J, et al. Subcortical brain volume abnormalities in 2028 individuals with schizophrenia and 2540 healthy controls via the ENIGMA consortium. *Mol Psychiatry* 2016; 21:547-53.
53. Hibar DP, Westlye L, van Erp T, et al. Subcortical volumetric abnormalities in bipolar disorder. *Mol Psychiatry* 2016;doi:10.1038/mp.2015.227.
54. Morey RA, Petty CM, Xu Y, et al. A comparison of automated segmentation and manual tracing for quantifying hippocampal and amygdala volumes. *Neuroimage* 2009;45:855-66.
55. Mulder ER, de Jong RA, Knol DL, et al. Hippocampal volume change measurement: quantitative assessment of the reproducibility of expert manual outlining and the automated methods FreeSurfer and FIRST. *Neuroimage* 2014;92:169-81.
56. Blake DD, Weathers FW, Nagy LM, et al. The development of a Clinician-Administered PTSD Scale. *J Trauma Stress* 1995;8:75-90.
57. Wechsler D. Wechsler Test of Adult Reading: WTAR. Psychological Corporation; 2001.
58. Sapolsky RM. Glucocorticoids and hippocampal atrophy in neuropsychiatric disorders. *Arch Gen Psychiatry* 2000;57:925-35.
59. Moghaddam B, Bolinao ML, Stein-Behrens B, et al. Glucocorticoids mediate the stress-induced extracellular accumulation of glutamate. *Brain Res* 1994;655:251-4.
60. Bailey CH, Giustetto M, Huang Y-Y, et al. Is heterosynaptic modulation essential for stabilizing hebbian plasticity and memory. *Nat Rev Neurosci* 2000;1:11-20.
61. O'Carroll CM, Morris RG. Heterosynaptic co-activation of glutamatergic and dopaminergic afferents is required to induce persistent long-term potentiation. *Neuropharmacology* 2004;47:324-32.
62. Bruel-Jungerman E, Davis S, Rampon C, et al. Long-term potentiation enhances neurogenesis in the adult dentate gyrus. *J Neurosci* 2006; 26:5888-93.
63. Rocher C, Spedding M, Munoz C, et al. Acute stress-induced changes in hippocampal/prefrontal circuits in rats: effects of antidepressants. *Cereb Cortex* 2004;14:224-9.
64. Jay TM, Rocher C, Hote M, et al. Plasticity at hippocampal to prefrontal cortex synapses is impaired by loss of dopamine and stress: importance for psychiatric diseases. *Neurotox Res* 2004;6:233-44.
65. Caudle WM, Richardson JR, Wang MZ, et al. Reduced vesicular storage of dopamine causes progressive nigrostriatal neurodegeneration. *J Neurosci* 2007;27:8138-48.
66. Bertolino A, Rubino V, Sambataro F, et al. Prefrontal-hippocampal coupling during memory processing is modulated by COMT Val158Met genotype. *Biol Psychiatry* 2006;60:1250-8.
67. Schott BH, Seidenbecher CI, Fenker DB, et al. The dopaminergic midbrain participates in human episodic memory formation: evidence from genetic imaging. *J Neurosci* 2006;26:1407-17.
68. Schulz-Heik RJ, Schaefer M, Eliez S, et al. Catechol-O-methyltransferase Val158Met polymorphism moderates anterior cingulate volume in posttraumatic stress disorder. *Biol Psychiatry* 2011;70:1091-6.
69. Lisman JE, Grace AA. The hippocampal-VTA loop: controlling the entry of information into long-term memory. *Neuron* 2005;46:703-13.
70. Menezes J, Alves N, Borges S, et al. Facilitation of fear extinction by novelty depends on dopamine acting on D1-subtype dopamine receptors in hippocampus. *Proc Natl Acad Sci U S A* 2015;112:E1652-8.
71. Cha J, Carlson JM, DeDora DJ, et al. Hyper-reactive human ventral tegmental area and aberrant mesocorticolimbic connectivity in overgeneralization of fear in generalized anxiety disorder. *J Neurosci* 2014;34:5855-60.
72. Lee TM, Yip JT, Jones-Gotman M. Memory deficits after resection from left or right anterior temporal lobe in humans: a meta-analytic review. *Epilepsia* 2002;43:283-91.
73. Uddo M, Vasterling JJ, Brailey K, et al. Memory and attention in combat-related post-traumatic stress disorder (PTSD). *J Psychopathol Behav Assess* 1993;15:43-52.

Correction: Ventral striatal hypoactivation is associated with apathy but not diminished expression in patients with schizophrenia

In the article "Ventral striatal hypoactivation is associated with apathy but not diminished expression in patients with schizophrenia" by Kirschner and colleagues¹ published in the May 2016 issue of the journal, the paragraph on "experimental design and task" in the Methods section should have mentioned that the task variant used was originally developed by Simon and colleagues.² The text and the reference list have been updated in the online version of the article, which is available at jpn.ca. We apologize for the error.

Reference

1. Kirschner M, Hager OM, Bischof M, et al. Ventral striatal hypoactivation is associated with apathy but not diminished expression in patients with schizophrenia. *J Psychiatry Neurosci* 2016;41(3):152-161.
2. Simon JJ, Cordeiro SA, Weber M-A, et al. Reward system dysfunction as a neural substrate of symptom expression across the general population and patients with schizophrenia. *Schizophr Bull* 2015;41:1370-8.