

Cardiorespiratory concerns shape brain responses during automatic panic-related scene processing in patients with panic disorder

Katharina Feldker, Dr. rer. nat; Carina Yvonne Heitmann, Dr. rer. nat;
Paula Neumeister, Dr. rer. nat; Leonie Brinkmann, Dr. rer. nat;
Maximilian Bruchmann, Dr. rer. nat; Pienie Zwitserlood, PhD; Thomas Straube, PhD

Background: Increased automatic processing of threat-related stimuli has been proposed as a key element in panic disorder. Little is known about the neural basis of automatic processing, in particular to task-irrelevant, panic-related, ecologically valid stimuli, or about the association between brain activation and symptomatology in patients with panic disorder. **Methods:** The present event-related functional MRI (fMRI) study compared brain responses to task-irrelevant, panic-related and neutral visual stimuli in medication-free patients with panic disorder and healthy controls. Panic-related and neutral scenes were presented while participants performed a spatially non-overlapping bar orientation task. Correlation analyses investigated the association between brain responses and panic-related aspects of symptomatology, measured using the Anxiety Sensitivity Index (ASI). **Results:** We included 26 patients with panic disorder and 26 healthy controls in our analysis. Compared with controls, patients with panic disorder showed elevated activation in the amygdala, brainstem, thalamus, insula, anterior cingulate cortex and midcingulate cortex in response to panic-related versus neutral task-irrelevant stimuli. Furthermore, fear of cardiovascular symptoms (a subcomponent of the ASI) was associated with insula activation, whereas fear of respiratory symptoms was associated with brainstem hyperactivation in patients with panic disorder. **Limitations:** The additional implementation of measures of autonomic activation, such as pupil diameter, heart rate, or electrodermal activity, would have been informative during the fMRI scan as well as during the rating procedure. **Conclusion:** Results reveal a neural network involved in the processing of panic-related distractor stimuli in patients with panic disorder and suggest an automatic weighting of panic-related information depending on the magnitude of cardiovascular and respiratory symptoms. Insula and brainstem activations show function-related associations with specific components of panic symptomatology.

Introduction

Recurring sudden panic attacks and anxious apprehension are the 2 core symptoms for diagnosing panic disorder.¹ Panic-related anxious apprehension is characterized by worry about future attacks and by fear of bodily and cognitive sensations of anxiety.² In patients with panic disorder and other anxiety disorders, threat-related stimuli, which are by definition salient, activate neural mechanisms that facilitate fast and preferred processing.³⁻⁶ This automatic attention to threat-related stimuli is associated with detection and evaluation mechanisms, reorientation of resources and amplification of processing mechanisms.⁷

Reviews of functional imaging studies propose a network including the amygdala, brainstem, thalamus, insula, anterior

cingulate cortex (ACC), midcingulate cortex (MCC) and medial prefrontal cortex (mPFC) as neural underpinnings of altered threat processing in patients with panic disorder.⁸⁻¹⁰ However, the neural basis of the processing of panic-related stimuli when these are irrelevant for the task at hand has not commonly been investigated in patients with panic disorder. Using an emotional Stroop task, greater activation in patients with panic disorder than in healthy controls was found in the prefrontal cortex,¹¹ ACC, thalamus, amygdala, hippocampus and inferior parietal cortex.⁵ With spatially overlapping emotional word-face pairs that required a response to faces only, Chechko and colleagues¹² reported decreased ACC and frontal gyrus activation, but increased activation in the amygdala and brainstem in patients with remitted panic disorder for emotionally incongruent compared to congruent pairs.

Correspondence to: K. Feldker, Institute of Medical Psychology and Systems Neuroscience, Von-Esmarch-Straße 52, D-48149 Muenster, Germany; katharina.feldker@uni-muenster.de

Submitted Nov. 24, 2017; Revised Mar. 30, 2017; Revised Apr. 28, 2017; Accepted May 1, 2017; Online first Sept. 26, 2017

DOI: 10.1503/jpn.160226

However, studies investigating neural responses to task-irrelevant, ecologically valid panic-related stimuli under experimental conditions in which attention to distracting emotional information is not required at all hardly exist.

Moreover, and most importantly, it remains to be specified how brain responses under these conditions are associated with symptoms of panic disorder. Patients with this disorder have varied anxiety sensitivity, defined as the fear of anxiety-related bodily sensations derived from beliefs that these symptoms have harmful physical, psychological, or social consequences.^{13,14} The Anxiety Sensitivity Index (ASI) has been shown to effectively measure fear for physical, psychological, or social aspects associated with the experience of anxiety.^{15,16} Anxiety severity is associated with the development of spontaneous panic attacks in healthy young adults, and fear of physical concerns (a subcomponent of anxiety severity) plays a particularly important role in predicting the course of panic disorder.^{17–20} A better understanding of the role of anxiety severity in panic disorder could thus help to design prevention programs, specify therapeutic interventions and might further enrich diagnostic processes. Poletti and colleagues²¹ reported positive correlations of anxiety severity with neural activation in the PFC, ACC and insula in patients with panic disorder. Thus, preliminary behavioural and neural data underscore the need for further research into the role of anxiety severity subconstructs in patients with the disorder. It is unknown whether anxiety severity modulates neural effects of automatic, panic-related stimulus processing. Research on anxiety severity has led to revisions in factor structure, currently suggesting 4 subscales (fear of cardiovascular symptoms, fear of respiratory symptoms, fear of loss of control, fear of publicly observable symptoms) in the German ASI-4.¹⁵ The ASI-4 factor structure that describes fear of physical symptoms in more detail allows us to disentangle the predictive value of fear of cardiovascular and respiratory symptoms.¹⁵

The present event-related functional MRI (fMRI) study aimed to elucidate the neural underpinnings of processing panic-related stimuli in patients with panic disorder when these stimuli are spatially separate from the targets and completely task-irrelevant.⁷ We used a recently developed standardized set of panic-related and neutral scenes as distractors while participants had to solve a concurrent but spatially nonoverlapping bar discrimination task. We further addressed how neural effects are shaped by panic-related aspects of anxiety sensitivity measured with the ASI-4, using correlation analysis.

Methods

Participants

We recruited patients with panic disorder and healthy controls through public advertisement and in collaboration with an outpatient clinic. Psychiatric medication, neurologic disorders, and fMRI contraindications served as exclusion criteria for both groups. Psychotic or bipolar disorders, suicidal ideations, and drug dependence or abuse within the last 10 years were additional exclusion criteria for the panic dis-

order group. Controls also had to be free of any psychiatric diagnosis within the last 10 years. Patients had to be free of psychiatric medication for at least 3 months and had to refrain from “as needed” medication for at least 1 week before the fMRI session. An experienced clinical psychologist interviewed all patients and controls using the Structured Clinical Interview for DSM-IV Axis I Disorders (SCID). Participants completed questionnaires on demographic and clinical data. The German ASI-4,¹⁵ a 24-item questionnaire, was administered to assess fear of anxiety-related symptoms on fear of cardiovascular symptoms, respiratory symptoms, loss of control and publicly observable symptoms using a 5-point Likert scale (0 = don’t agree at all; 4 = agree completely). All participants gave written informed consent. The study was approved by the ethics committee of the University of Muenster and conforms with the Declaration of Helsinki.

Stimuli

We used the Panic-related Picture Set Münster (PAPS-M), comprising 50 panic-related and 50 neutral scenes. The set was developed in a 3-step pilot study, including an extensive Web search, a clinical expert rating and a patient rating. Panic-related scenes display symptoms related to panic attacks (e.g., shortness of breath, hyperventilation, heart palpitations, chest pain, trembling or shaking, dizziness, fainting), fears (e.g., having a heart attack) and agoraphobia-related situations (e.g., crowded bus, dark tunnel, glass elevator).²²

Procedure

During the 6 minute 56 second experimental stimulation, each of the 50 panic-related and 50 neutral distractor scenes was presented once, centred on the screen, in an event-related design. Two circles (target) with a small line inside were presented above and below the concurrently displayed scene (distractor), and participants performed a bar orientation task (Fig. 1; modified from Straube and colleagues²³ and Wiens and colleagues²⁴). Lines were either vertical or horizontal and had either the same or different orientation in both circles (target stimuli). Participants indicated by button press with their right index and middle fingers whether orientation was the same or different in both circles. Orientation was the same on half of the trials and different for the other half. Trials were classified as errors when participants gave an incorrect response, missed a trial or pressed both buttons. This task can be categorized as a concurrent but distinct target–distractor task (CDTD) or directed-attention task, which is well suited to investigate distractor-driven attentional processes.^{7,25} Each scene (distractor) and the 2 circles (target) were presented for 800 ms, separated by a white fixation cross (1280–12200 ms, mean 3360 ms). Neutral and panic-related scenes were presented in random sequence and optimized and counterbalanced with the Optseq algorithm (<https://surfer.nmr.mgh.harvard.edu/optseq/>), which implements temporal jitter to increase signal discriminability.²⁶ Stimuli were rear-projected onto a screen that the participants viewed through a mirror on the MRI head coil.

Participants rated all 100 scenes within 7 days after the fMRI experiment in a postscanning rating session. After presentation of a 2 s stimulus, participants rated the scenes with regard to valence, arousal and anxiety. A 9-point Likert scale was used to assess valence (1 = very unpleasant; 5 = neutral; 9 = very pleasant), arousal (1 = not arousing; 9 = very arousing) and anxiety (1 = not anxiety-inducing; 9 = very anxiety-inducing).

Behavioural data analysis

Trials with premature button presses (< 300 ms), delayed responses (+2 standard deviations from the individual log-transformed mean) and erroneous button presses were discarded (315 trials, 6% of all data). To compensate for a skewed reaction time distribution, all reaction time analyses were performed on log-transformed data. Reaction times and rating data were analyzed by means of 2×2 repeated-measures analyses of variance (ANOVA) with emotion (panic-related, neutral) as a within-subjects factor and group (panic disorder,

healthy control) as a between-subjects factor. We considered results to be significant at $p < 0.05$, and post hoc comparisons were Bonferroni-corrected for multiple testing.

fMRI acquisition and analysis

We recorded structural brain information and blood oxygen level-dependent (BOLD) responses using a Magnetom PRISMA 3 T MRI scanner with a 20-channel head matrix coil (Siemens Medical Solutions). We recorded a T_1 -weighted MPRAGE structural volume with 192 slices for anatomic localization. We conducted a run of 225 volumes using a T_2^* -weighted echo-planar sequence with the following parameters: echo time (TE) 30 ms, flip angle 90° , matrix 92×92 voxels, field of view (FOV) 208 mm, repetition time (TR) 2080 ms. Each volume consisted of 36 axial slices (thickness 3 mm, gap 0.3 mm, in-plane resolution 2.26×2.26 mm²). The volumes were tilted approximately 20° clockwise from the anterior-posterior commissure line, to minimize susceptibility artifacts




Fig. 1: Schematic overview of a trial in the concurrent but distinct target-distractor task.

in inferior parts of anterior brain areas. A shimming field was applied before functional imaging to minimize external magnetic field inhomogeneities.

We conducted our fMRI data using BrainVoyager QX software version 2.4 (Brain Innovation). The first 10 volumes of each run were discarded from analysis to ensure steady state tissue magnetization. After the last trial, the fixation cross was presented for an additional 15 volumes to allow for data acquisition in randomizations with a very short last interstimulus interval.²⁶ All volumes were realigned to the first volume to minimize artifacts due to head movements and were then resampled to a voxel size of $2 \times 2 \times 2 \text{ mm}^3$. Slice-time correction and spatial (6 mm full-width at half-maximum [FWHM] isotropic Gaussian kernel) and temporal smoothing (high pass filter: 10 cycles in time course [0.023 Hz]; low pass filter: 2.8 s FWHM Gaussian kernel; linear trend removal) were applied. The anatomic and functional images were coregistered and normalized to Talairach space.²⁷ We used the normalization procedure implemented by default in BrainVoyager.

Small volume-corrected and whole-brain analyses

For statistical analyses, we calculated multiple linear regressions modelling the signal time course at each voxel. The expected BOLD signal change for each predictor was modelled with a canonical double γ hemodynamic response function (HRF). Predictors of interest were panic-related and neutral scenes. Only trials in which participants gave a correct answer were included in the analysis. Analogous to reaction-time data analysis, trials including errors were modulated in a separate predictor. This predictor and 6 motion parameters (to account for movement artifacts) were included as regressors of no interest into the model. First, we generated voxel-wise statistical maps and computed percent-standardized predictor estimates (b weights) for each participant. We analyzed predictor estimates across participants by means of *t* tests in specific regions of interest (ROIs).

Based on the current panic disorder imaging literature, the brainstem, amygdala, insula, thalamus, ACC, MCC, and mPFC served as ROIs.^{8,9} The ROIs (amygdala, insula, thalamus, mPFC, ACC and MCC) were created based on the Automated Anatomic Labelling (AAL) atlas included in the Wake Forest University (WFU) pick atlas^{28,29} with a 1 mm dilation factor for the amygdala and insula. For the mPFC, we used the 2 AAL templates "Frontal_Med_Orb" and "Frontal_Sup_Medial." The ROIs for the brainstem were downloaded from the digitized version of the Talairach atlas (www.talairach.org/nii/gzip/). We converted obtained Montreal Neurological Institute (MNI) coordinates to Talairach space using MATLAB version 8.2 (MathWorks) using the ICBM-152 routine proposed by Lancaster and colleagues.³⁰ Peak voxel labelling was supported by Talairach Dameon software 31 and verified using the Mai atlas.³²

For statistical analyses, we used a cluster-based permutation (CBP) approach.^{33,34} This approach requires no assumptions about the test statistic distribution and has recently been shown to be more valid than classical parametric fMRI analyses and to offer precise control of the false-discovery

rate.³⁴ All permutation tests were performed with 1000 permutations.^{35,36} For each permutation, the individual b maps (panic-related – neutral) were randomly assigned without replacement to 1 of the 2 experimental groups. Voxel threshold was set at $p_{\text{voxel}} < 0.005$ to balance between type I and type II error. We calculated cluster mass by adding all *F* values in neighbouring significant voxels. We compared the cluster mass observed in the contrast of interests with the distribution of the maximal cluster mass observed in each of the 1000 permutations. Cluster masses larger or equal to the 95th percentile of the permutation distribution were considered to be statistically significant clusters (i.e., $p_{\text{cluster}} < 0.05$).

We performed correlational analyses for ASI subscale scores (fear of cardiovascular symptoms, fear of respiratory symptoms, fear of publicly observable symptoms, fear of loss of control) to investigate the moderating influence of these scores on neural responses to panic-related versus neutral stimuli in patients with panic disorder. Using the CBP approach, individual b maps (panic-related > neutral) of anatomic regions in which differential between-group effects were detected were randomly assigned to the individual questionnaire subscale scores. The cluster mass was then calculated by adding all correlation coefficients of neighbouring significant voxels. We considered clusters with $p_{\text{cluster}} < 0.0125$ (corrected for number of ASI subscales) to be statistically significant.

Results

Participants

Of the 29 patients with panic disorder originally recruited, 1 aborted the scanning session owing to panic symptoms, and we discarded the data sets of 2 participants for whom 20 trials were classified as errors. This left a final sample of 26 patients with panic disorder (age range 18–46 yr) and 26 controls (age range 19–32 yr) comparable for age, sex and education. The demographic and clinical characteristics of the study groups are shown in Table 1. All participants were native German speakers, had normal or corrected-to-normal vision, and were right-handed, as assessed using the Edinburgh Handedness Inventory. Thirteen of the 26 patients had a primary diagnosis of panic disorder (DSM 300.01) and the other 13 had a primary diagnosis of panic disorder with agoraphobia (DSM 300.21). Six of the 26 patients were undergoing psychotherapy at the time of study participation.

Behavioural data

Patients with panic disorder rated scenes as more unpleasant, more arousing and more anxiety-inducing than healthy controls. Both groups rated panic-related stimuli as more unpleasant, more arousing and more anxiety-inducing than neutral scenes (Fig. 2 and Appendix 1, available at jpn.ca/160226-a1). Arousal and anxiety levels were significantly higher in patients with panic disorder than in controls for panic-related versus neutral scenes.

In the final sample, on average 94 of 100 (range 87–98) trials per person were included. The number of trials excluded

owing to errors or outliers did not differ between groups (average number of excluded trials: 5 in controls v. 7 in patients with panic disorder; $t_{50} = 1.47$, $p = 0.15$). Analysis of log-transformed reaction times showed neither significant main effects (emotion: $F_{1,50} = 0.083$; group: $F_{1,50} = 0.669$) nor a significant group \times emotion interaction ($F_{1,50} = 0.767$, all $p > 0.1$; reaction times in the panic disorder group: disorder-related = $900.22 \text{ ms} \pm 96.67 \text{ ms}$, neutral = $894.32 \text{ ms} \pm 83.45 \text{ ms}$; reaction times in healthy controls: disorder-related = $876.02 \text{ ms} \pm 97.71 \text{ ms}$, neutral = $878.29 \text{ ms} \pm 94.83 \text{ ms}$).

fMRI data

Small volume-corrected analysis

Patients with panic disorder showed stronger emotion effects (always referring to panic-related > neutral scenes) than healthy controls in the amygdala, brainstem, insula, thalamus (lateral nucleus), ACC and MCC (Table 2 and Fig. 3). In the mPFC, activation in response to panic-related versus neutral scenes did not differ between patients with panic disorder and controls. No hyperactivations in controls versus patients for panic-related versus neutral scenes were observed.

Whole-brain analysis

Whole-brain analysis yielded 2 clusters of greater activation in patients with panic disorder versus controls for panic-related versus neutral scenes. The first cluster encompassed brainstem regions, as shown in small volume-corrected analysis,

and extended to cerebellar and posterior cingulate cortex (PCC) regions. The second cluster encompassed the insula effect and extended to inferior frontal and temporal regions (Table 3). Healthy controls showed no clusters of activation




Fig. 2: Mean postscanning scene ratings for valence, arousal and anxiety (disorder-related – neutral) for patients with panic disorder and healthy controls. Ratings were given on 9-point Likert scales as follows: valence, 1 = negative, 5 = neutral, 9 = positive; arousal, 1 = calm, 9 = intense; anxiety, 1 = low, 9 = high.

Table 1: Demographic and clinical characterization of patients with panic disorder and healthy controls

Characteristic	Group; mean \pm SD*		Statistic	p value
	Panic disorder	Control		
Sex, female:male	20:6	19:7	$\chi^2 = 0.103$	0.75
Age, (range) yr	24.88 ± 6.12 (18–46)	24.46 ± 2.79 (19–32)	$t = -0.321$	0.75
Education, yr	12.46 ± 0.99	12.52 ± 1.05	$t = 0.205$	0.84
Diagnosis, no.	—	—	—	—
Panic disorder	13	NA	—	—
Panic disorder with agoraphobia	13	NA	—	—
PAS score	20.46 ± 7.05	NA	—	—
ASI score				
Total	45.92 ± 2.88	10.12 ± 1.67	$t = -10.76$	< 0.001
Cardiovascular symptoms	9.58 ± 0.98	1.5 ± 0.40	$t = -7.63$	< 0.001
Respiratory symptoms	14.23 ± 1.25	2.69 ± 0.60	$t = -8.31$	< 0.001
Publicly observable symptoms	11.92 ± 1.03	4.65 ± 0.56	$t = -6.20$	< 0.001
Loss of control	10.19 ± 0.90	1.27 ± 0.39	$t = -9.10$	< 0.001
BDI score	14.23 ± 8.94	0.92 ± 2.08	$t = -7.39$	< 0.001
Comorbidities, no.				
Mild depressive episode	3	—	—	—
Generalized anxiety disorder	4	—	—	—
Somatization disorder and hypochondria	1	—	—	—
Social or specific phobia	2	—	—	—
Bulimia nervosa	1	—	—	—
OCD	1	—	—	—


ASI = Anxiety Sensitivity Scale; BDI = Beck Depression Inventory; NA = not applicable; OCD = obsessive-compulsive disorder; PAS = Panic and Agoraphobia Scale; SD = standard deviation.

*Unless indicated otherwise.

Table 2: Significant hyperactivations for disorder-related versus neutral scenes across all patients relative to healthy controls revealed by small volume–corrected analysis ($p \leq 0.05$, corrected)

Region	Lateralization	Talairach coordinates of peak voxel (x, y, z)	t maximum	t average	k
Amygdala	L	-17, -7, -8	3.49	3.196	17
Amygdala	R	23, 3, -14	3.284	3.059	15
Insula	L	-31, -1, 18	4.166	3.313	45
Insula	L	-45, 8, -8	4.365	3.298	84
Brainstem	L	-3, -35, -8	4.311	3.444	134
Brainstem	L	-7, -33, -28	4.079	3.431	37
Thalamus, lat part	L	-15, -19, 6	4.548	3.433	41
Anterior cingulate cortex	L	-3, 19, 34	3.926	3.314	21
Anterior cingulate cortex	L	-3, 37, 16	3.415	3.108	22
Midcingulate cortex	L/R	-1, -1, 30	3.868	3.282	100
Midcingulate cortex	L/R	-5, 19, 38	4.26	3.412	60

L = left; R = right.

**Fig. 3:** Differential brain activation for disorder-related compared with neutral distractor scenes in patients with panic disorder versus healthy controls in a priori–defined regions of interest (panic disorder > control, panic-related > neutral, all $p < 0.005$, uncorrected; $p < 0.05$, corrected; only selected effects are shown). ACC = anterior cingulate cortex; L = left; MCC = midcingulate cortex; R = right.

higher than patients with panic disorder for the contrast panic-related versus neutral stimuli.

The investigation of the influence of ASI-4 subscales on differential brain activation in patients with panic disorder resulted in a significant positive correlation between fear of cardiovascular symptoms and anterior insula activation (peak voxel: $x, y, z = -41, 19, -5$; $\max R = 0.718$, $\text{avg} R = 0.557$, $k = 246$ voxels, $p = 0.003$) and between fear of respiratory symptoms and brainstem activation (peak voxel: $x, y, z = -12, -24, -24$; $\max R = 0.692$, $\text{avg} R = 0.563$, $k = 136$ voxels, $p = 0.004$; Fig. 4).

Discussion

The present study investigated in patients with panic disorder the neural correlates of automatic processing of panic-related visual scenes that were task-irrelevant and spatially distinct from task targets. Comparing panic-related versus neutral scenes, patients with panic disorder showed greater activation than controls in the amygdala, brainstem, thalamus, insula, ACC and MCC, whereas behavioural responses did not differ between the groups. Fear of cardiovascular symptoms was the strongest predictor of emotion effects (always referring to panic-related > neutral) in the insula, whereas fear of respiratory symptoms predicted emotion effects in the brainstem in patients with panic disorder.

Next to the general hyperactivation in several brain regions in patients with panic disorder during processing of task-irrelevant, panic-related stimuli, activation in 2 key regions — the insula and brainstem — was modulated by specific aspects of panic symptomatology. The insula, associated with interoceptive processing and the processing of one's own bodily symptoms, plays an important role in panic disorder.^{9,37,38} Hyperactivation in the anterior insula in our patients is best explained by fear of cardiovascular symptoms. Earlier studies found anxiety severity to correlate positively with insula hyperactivation in response to masked fearful faces in healthy controls and with insula activation in patients with panic disorder during emotional processing of facial affect expressions.^{21,39} Killgore and colleagues³⁹ empha-

sized the role of specific subscales of anxiety severity and reported fear of physical symptoms to significantly correlate with anterior insular activation. Our findings of insular activation, as driven by fear of cardiovascular symptoms, match this result and extend it to a clinical sample that is characterized by its enhanced sensibility to bodily symptoms relative to other anxiety disorders.⁴⁰ The insula hyperactivity of patients with panic disorder when processing disorder-related scenes may thus be specifically associated with their sensibility to and monitoring of bodily symptoms of anxiety.^{41–43} Findings suggest that those who fear physical symptoms more also pay them more attention, and vice versa. Notably, we observed greater panic-related insula activation despite panic-related scenes requiring no explicit processing, which indicates a low threshold for interoceptive processing in patients with panic disorder — a marker that might present an etiological or maintaining factor in panic disorder.

Furthermore, our results showed that task-irrelevant panic-related scenes trigger greater brainstem activation in patients with panic disorder than in healthy controls (panic-related > neutral), and fear of respiratory symptoms predicted differential brainstem activation in patients with panic disorder. Brainstem alterations in patients with panic disorder have been reported in anatomic^{44–46} and functional imaging studies.^{12,47–49} Aberrant activation in the brainstem, a site of homeostatic integration, may be closely linked to the changes in chemoreception and cardiorespiratory control perceived by patients with panic disorder.^{10,50–52} Using a carbon dioxide challenge, Goossens and colleagues⁵³ reported increased brainstem activation in response to hypercapnia in patients with panic disorder compared with healthy controls. Although limited resolution does not allow for specific localization of subregions, coordinates suggest an involvement of pontine nuclei and locus coeruleus, a carbon dioxide/H⁺-sensitive brain site involved in communication of respiratory- and stress-induced activation changes.^{54,55} The present study adds to what is known about the association between the brainstem and respiratory symptoms by linking increased brainstem activation to the subjective fear of respiratory

Table 3: Significant hyperactivations for disorder-related versus neutral scenes across all patients relative to healthy controls revealed by whole-brain analysis in 2 clusters ($p \leq 0.05$, corrected)*

Region	Lateralization	Talairach coordinates of peak voxel (x, y, z)	F maximum	F average	k
Cluster 1	L/R	1, -39, -2	4.38	3.40	504
Culmen	R	1, -39, -2	4.38	3.40	378
Posterior cingulate cortex	R	17, -43, 8	4.20	3.48	68
Brainstem/culmen	L	-10, -30, -7	4.03	3.37	19
Posterior cingulate cortex	R	24, -54, 10	3.55	3.16	39
Cluster 2	L	-47, 8, -3	4.51	3.28	399
Superior temporal gyrus	L	-47, 8, -3	4.51	3.26	185
Insula	L	-38, -1, -7	4.11	3.34	76
Precentral gyrus	L	-61, 5, 13	4.01	3.24	69
Temporal lobe	L	-35, -8, -13	3.88	3.36	30
Inferior frontal gyrus	L	-48, 3, 17	3.74	3.25	39

L = left ; R = right.

*The watershed algorithm of NeuroElf (v0.9c; <http://neuroelf.net/>); i.e., the splitclustercoords function) was used to assess local maxima of clusters.

symptoms. In his “false suffocation alarms” hypothesis, Klein⁵⁶ postulated that patients with panic disorder have a pathologically altered suffocation alarm monitor that results in a carbon dioxide hypersensitivity, forming the basis for sudden panic attacks. Integrating findings by Goossens and colleagues⁵³ as well as the present findings, brainstem hyperactivation and its association with the subjective fear of respiratory symptoms might present a neurobiological substrate of such a monitoring system. Our results thus support the idea of increased brainstem activation in patients with panic disorder as a marker of an oversensitive alarm system that might predispose patients to the development of panic attacks. Replication of results, a more detailed postscanning interview on feelings of suffocation, a measurement of end-tidal carbon dioxide as well as more fine-grained analysis of brainstem subregional findings are necessary to further explore the neural basis of the false suffocation alarm hypothesis.

Several further ROIs, such as the amygdala, thalamus, ACC and MCC, showed increased activation in response to threat in patients with panic disorder, but without an association with symptom scores. Amygdala hyperactivation in patients with panic disorder has been interpreted as hyperresponsivity to environmental cues, eliciting full-scale threat-related responses.⁵⁷ Remarkably, findings of greater amygdala activation in patients with panic disorder than in

healthy controls are inconsistent,^{9,58} and panic attacks have even been observed in patients with amygdala lesions.^{59–61} Thus, the amygdala seems to be associated with relevance detection and salience processing,^{62,63} initiating a cascade of panic-related activations without being necessarily involved in (all) full-blown panic attacks.⁶⁰ The thalamus, as a sensory relay station, is important for the fast processing of incoming visual input and weighing its relevance. Its activation is modulated by selective attention, which gains special significance in the present study owing to the task-irrelevance of panic-related stimuli.^{64,65}

For the resolution of conflict caused by task-irrelevant panic-related stimuli, which closely links emotion and cognition, the cingulate cortex has been proposed as a major neural structure.^{66,67} The dorsal cingulate cortex plays an important role in overcoming interference due to emotional distraction,⁶⁸ which is relevant to understanding the co-occurrence of elevated emotion effects in the fear network of patients with panic disorder in the absence of significant differences on the behavioural level. Heightened cingulate cortex activation has also been linked to enhanced sensory sensitivity, including exaggerated scanning for threats (i.e., hypervigilance).^{66,69} The dorsal anterior cingulate cortex has been linked to the active monitoring of emotions (e.g., the appraisal of an emotion-inducing stimulus), whereas the ventral cingulate cortex has




Fig. 4: Correlation of panic disorder parameter estimates in the left insula (panic-related > neutral) and fear of cardiovascular symptoms as well as correlation of panic disorder parameter estimates in the brainstem (panic-related > neutral) and fear of respiratory symptoms. L = left.

mainly been associated with emotion regulation.⁷⁰ With regard to the present data, this suggests that patients with panic disorder might consciously attribute threat to the disorder-related stimuli, but lack the capacity to activate emotion regulation processes associated with more ventral cingulate cortex activations. According to Seeley and colleagues,⁶ the dorsal cingulate cortex represents the key node in the salience network that reflects paralimbic emotional salience processing and is modulated by subjective anxiety ratings.⁶

The whole-brain analysis showed larger effects in response to panic-related stimuli in patients with panic disorder than in healthy controls in a brainstem/cerebellum/PCC cluster and an anterior insula/inferior-frontotemporal cluster. The hyperactivation for panic-related scenes in visual processing regions in patients with panic disorder underlines the anxiety-related relevance of these scenes for patients, given that the affective significance of stimuli enhances their visual processing.⁷¹⁻⁷³ Emotional salience and simultaneous attentional influences have been observed to modulate sensory processing.⁷⁴

Limitations

Some limitations of our study need to be considered. Although the sample of 26 patients can be considered large in the field of clinical affective neuroscience, greater sample sizes are needed to increase power in statistical analyses. Six of the 26 patients with panic disorder were undergoing psychotherapy at the time of study participation; however, the Panic and Agoraphobia Scale scores of these 6 patients indicated mild to moderately severe impairment. We implemented a recently developed set of complex visual disorder-related stimuli. However, panic disorder is characterized by strong sensations, including multiple sensory modalities. Thus, anticipation of disorder-related stimuli on other sensory modalities should be compared with or combined with visual emotional processing in future studies. In addition, future studies should also include general threatening stimuli in order to disentangle the effects of generally anxiety-inducing and panic-related stimuli. The additional implementation of measures of autonomic activation, such as pupil diameter, heart rate, or electrodermal activity, would have been informative during the fMRI scan as well as during the rating procedure. More precisely, a concurrent measurement of autonomic measures might help us gain further insight into the involvement of brainstem substructures, such as the locus coeruleus. Future studies should implement such measures to further investigate how subjective and objective measures from anxiety-related body systems translate to the brain.

Conclusion

Taken together, the present study provides new insights about neural correlates of automatic processing and their modulation by key aspects of symptomatology in patients with panic disorder. The disorder can be described as the subjective sense of danger to the core survival systems, such as respiration and heartbeat. Findings in the insula and brainstem suggest that fear of cardiovascular and fear of respira-

tory symptoms represent core features of panic disorder and are associated with neural activation in regions that are important for monitoring body states and for signalling a potential state of alarm. The present study greatly adds to the knowledge on the neurobiological basis of these fears by revealing an association of subjectively reported panic disorder-specific fears and altered activation in the insula and brainstem in an automatic context.

Affiliations: From the Institute of Medical Psychology and Systems Neuroscience, University of Muenster, Muenster, Germany (Feldker, Heitmann, Neumeister, Brinkmann, Bruchmann, Straube); and the Institute for Psychology, University of Muenster, Muenster, Germany (Zwisterlood).

Funding: This work was supported by grants awarded to T. Straube by the German Research Society (Deutsche Forschungsgemeinschaft, (DFG/SFB/ TRR 58: C06, C07).

Competing interests: None declared.

Contributors: K. Feldker, C. Heitmann, P. Neumeister, P. Zwisterlood and T. Straube designed the study. K. Feldker, C. Heitmann and P. Neumeister acquired the data, which all authors analyzed. K. Feldker and L. Brinkmann wrote the article, which all authors reviewed and approved. All authors approved the final version to be published and can certify that no other individuals not listed as authors have made substantial contributions to the paper.

References

1. American Psychiatric Association. Diagnostic and statistical manual of mental disorders: DSM-IV-TR. 4th edition, text revision. Washington (DC): American Psychiatric Association; 2000.
2. Helbig-Lang S, Lang T, Petermann F, et al. Anticipatory anxiety as a function of panic attacks and panic-related self-efficacy: an ambulatory assessment study in panic disorder. *Behav Cogn Psychother* 2012;40:590-604.
3. Öhman A. The role of the amygdala in human fear: automatic detection of threat. *Psychoneuroendocrinology* 2005;30:953-8.
4. Reinecke A, Waldenmaier L, Cooper MJ, et al. Changes in automatic threat processing precede and predict clinical changes with exposure-based cognitive-behavior therapy for panic disorder. *Biol Psychiatry* 2013;73:1064-70.
5. van den Heuvel OA, Veltman D, Groenewegen H. Disorder-specific neuroanatomical correlates of attentional bias in obsessive-compulsive disorder, panic disorder, and hypochondriasis. *Arch Gen Psychiatry* 2005;62:922-33.
6. Seeley WW, Menon V, Schatzberg AF, et al. Dissociable intrinsic connectivity networks for salience processing and executive control. *J Neurosci* 2007;27:2349-56.
7. Carretié L. Exogenous (automatic) attention to emotional stimuli: a review. *Cogn Affect Behav Neurosci* 2014;14:1228-58.
8. de Carvalho MR, Dias GP, Cosci F, et al. Current findings of fMRI in panic disorder: contributions for the fear neurocircuitry and CBT effects. *Expert Rev Neurother* 2010;10:291-303.
9. Dresler T, Guhn A, Tupak SV, et al. Revise the revised? New dimensions of the neuroanatomical hypothesis of panic disorder. *J Neural Transm* 2013;120:3-29.
10. Gorman JM, Kent JM, Sullivan GM, et al. Neuroanatomical hypothesis of panic disorder, revised. *FOCUS J* 2004;2:426-39.
11. Dresler T, Attar CH, Spitzer C, et al. Neural correlates of the emotional Stroop task in panic disorder patients: an event-related fMRI study. *J Psychiatr Res* 2012;46:1627-34.

12. Chechko N, Wehrle R, Erhardt A, et al. Unstable prefrontal response to emotional conflict and activation of lower limbic structures and brainstem in remitted panic disorder. *PLoS One* 2009;4:e5537-11.
13. Onur E, Alkin T, Tural U. Panic disorder subtypes: further clinical differences. *Depress Anxiety* 2007;24:479-86.
14. Reiss S, McNally RJ. Expectancy model of fear. In: Reiss S, Bootzin RR (editors). *Theoretical issues in behavior therapy*. San Diego (CA): Academic Press; 1985:107-21.
15. Kemper CJ, Specht M, Volk S. Konstruktvalidität und Nutzen eines Verfahrens zur Erfassung der Angstsensitivität (Angstsensitivitätsindex-4) in einer Stichprobe von Patienten mit schlafbezogenen Atmungsstörungen. Bremen; 2010.
16. Reiss S. Expectancy model of fear, anxiety, and panic. *Clin Psychol Rev* 1991;11:141-53.
17. Hayward C, Killen JD, Kraemer HC, et al. Predictors of panic attacks in adolescents. *J Am Acad Child Adolesc Psychiatry* 2000;39:207-14.
18. McNally RJ. Anxiety sensitivity and panic disorder. *Biol Psychiatry* 2002;52:938-46.
19. Schmidt NB, Zvolensky MJ, Maner JK. Anxiety sensitivity: prospective prediction of panic attacks and Axis I pathology. *J Psychiatr Res* 2006;40:691-9.
20. Pilecki B, Arentoft A, McKay D. An evidence-based causal model of panic disorder. *J Anxiety Disord* 2011;25:381-8.
21. Poletti S, Radaelli D, Cucchi M, et al. Neural correlates of anxiety sensitivity in panic disorder: a functional magnetic resonance imaging study. *Psychiatry Res* 2015;233:95-101.
22. Feldker K, Heitmann CY, Neumeister P, et al. Brain responses to disorder-related visual threat in panic disorder. *Hum Brain Mapp* 2016;37:4439-53.
23. Straube T, Mentzel H-J, Miltner WHR. Neural mechanisms of automatic and direct processing of phobogenic stimuli in specific phobia. *Biol Psychiatry* 2006;59:162-70.
24. Wiens S, Syrjänen E. Directed attention reduces processing of emotional distracters irrespective of valence and arousal level. *Biol Psychol* 2013;94:44-54.
25. MacNamara A, Kappenman E, Black S, et al. Integrating behavioral and electrocortical measures of attentional bias toward threat. In: Caplovitz-Barrett K, Fox NA, Morgan GA, et al., editors. *Handbook of Self-Regulatory Processes in Development: New Directions and International Perspectives*. London (UK): Psychology Press; 2013.
26. Dale AM, Greve DN, Burock MA. Optimal stimulus sequences for event-related fMRI. Duesseldorf, Germany; 1999. Available: www.neurologie.uni-duesseldorf.de/HBM99/cd/methods/3095.html (accessed 2015 Sept. 4).
27. Talairach J, Tournoux P. *Co-planar stereotaxic atlas of the human brain*. New York (NY): Thieme; 1988.
28. Maldjian JA, Laurienti PJ, Kraft RA, et al. An automated method for neuroanatomic and cytoarchitectonic atlas-based interrogation of fMRI data sets. *Neuroimage* 2003;19:1233-9.
29. Tzourio-Mazoyer N, Landeau B, Papathanassiou D, et al. Automated anatomical labeling of activations in SPM using a macroscopic anatomical parcellation of the MNI MRI single-subject brain. *Neuroimage* 2002;15:273-89.
30. Lancaster JL, Tordesillas-Gutiérrez D, Martínez M, et al. Bias between MNI and Talairach coordinates analyzed using the ICBM-152 brain template. *Hum Brain Mapp* 2007;28:1194-205.
31. Lancaster JL, Woldorff MG, Parsons LM. Automated Talairach atlas labels for functional brain mapping. *Hum Brain Mapp* 2000;10:120-31.
32. Mai J, Assheur J, Paxinos G. *Atlas of the human brain*. 2. Ed. Amsterdam (Netherlands): Elsevier/Academic Press; 2004.
33. Kriegeskorte N, Simmons WK, Bellgowan PSF, et al. Circular analysis in systems neuroscience: the dangers of double dipping. *Nat Neurosci* 2009;12:535-40.
34. Eklund A, Nichols T, Knutsson H. Can parametric statistical methods be trusted for fMRI based group studies? [Internet]. Ithaca (NY): Cornell University Library; 2015. Available: <http://arxiv.org/abs/1511.01863> (accessed 2016 May 18).
35. Bullmore ET, Suckling J, Overmeyer S, et al. Global, voxel, and cluster tests, by theory and permutation, for a difference between two groups of structural MR images of the brain. *IEEE Trans Med Imaging* 1999;18:32-42.
36. Maris E, Oostenveld R. Nonparametric statistical testing of EEG- and MEG-data. *J Neurosci Methods* 2007;164:177-90.
37. Stern ER. Neural circuitry of interoception: new insights into anxiety and obsessive-compulsive disorders. *Curr Treat Options Psychiatry* 2014;1:235-47.
38. Khalsa SS, Lapidus RC. Can interoception improve the pragmatic search for biomarkers in psychiatry? *Front Psychiatry* 2016;7:121.
39. Killgore WDS, Britton JC, Price LM, et al. Neural correlates of anxiety sensitivity during masked presentation of affective faces. *Depress Anxiety* 2011;28:243-9.
40. Taylor S, Koch WJ, McNally RJ. How does anxiety sensitivity vary across the anxiety disorders? *J Anxiety Disord* 1992;6:249-59.
41. Craig AD. How do you feel — now? The anterior insula and human awareness. *Nat Rev Neurosci* 2009;10:59-70.
42. Holtz K, Pané-Farré CA, Wendt J, et al. Brain activation during anticipation of interoceptive threat. *Neuroimage* 2012;61:857-65.
43. Paulus MP, Stein MB. An insular view of anxiety. *Biol Psychiatry* 2006;60:383-7.
44. Fujiwara A, Yoshida T, Otsuka T, et al. Midbrain volume increase in patients with panic disorder. *Psychiatry Clin Neurosci* 2011;65:365-73.
45. Protopopescu X, Pan H, Tuescher O, et al. Increased brainstem volume in panic disorder: a voxel-based morphometric study. *Neuroreport* 2006;17:361-3.
46. Uchida RR, Del-Ben CM, Busatto GF, et al. Regional gray matter abnormalities in panic disorder: a voxel-based morphometry study. *Psychiatry Res Neuroimaging* 2008;163:21-9.
47. Lueken U, Straube B, Reinhardt I, et al. Altered top-down and bottom-up processing of fear conditioning in panic disorder with agoraphobia. *Psychol Med* 2014;44:381-94.
48. Tuescher O, Protopopescu X, Pan H, et al. Differential activity of rostral cingulate and brainstem in panic disorder and PTSD. *J Anxiety Disord* 2011;25:251-7.
49. Boshuisen ML, Ter Horst GJ, Paans AMJ, et al. rCBF differences between panic disorder patients and control subjects during anticipatory anxiety and rest. *Biol Psychiatry* 2002;52:126-35.
50. Pattinson KTS, Mitsis GD, Harvey AK, et al. Determination of the human brainstem respiratory control network and its cortical connections in vivo using functional and structural imaging. *Neuroimage* 2009;44:295-305.
51. Perna G, Guerriero G, Brambilla P, et al. Panic and the brainstem: clues from neuroimaging studies. *CNS Neurol Disord Drug Targets* 2014;13:1049-56.
52. Freire RC, Nardi AE. Panic disorder and the respiratory system: clinical subtype and challenge tests. *Rev Bras Psiquiatr* 2012;34:S32-41.
53. Goossens L, Leibold N, Peeters R, et al. Brainstem response to hypercapnia: a symptom provocation study into the pathophysiology of panic disorder. *J Psychopharmacol* 2014;28:449-56.
54. Esquivel G, Schruers KR, Maddock RJ, et al. Review: acids in the brain: A factor in panic? *J Psychopharmacol (Oxf)* 2010;24:639-47.
55. Itoi K, Sugimoto N. The brainstem noradrenergic systems in stress, anxiety and depression. *J Neuroendocrinol* 2010;22:355-61.
56. Klein DF. False suffocation alarms, spontaneous panics, and related conditions. An integrative hypothesis. *Arch Gen Psychiatry* 1993;50:306-17.

57. Rauch SL, Shin LM, Wright CI. Neuroimaging studies of amygdala function in anxiety disorders. *Ann N Y Acad Sci* 2003;985:389-410.
58. Etkin A, Wager TD. Functional neuroimaging of anxiety: a meta-analysis of emotional processing in PTSD, social anxiety disorder, and specific phobia. *Am J Psychiatry* 2007;164:1476-88.
59. Feinstein JS, Buzza C, Hurlmann R, et al. Fear and panic in humans with bilateral amygdala damage. *Nat Neurosci* 2013;16:270-2.
60. Khalsa SS, Feinstein JS, Li W, et al. Panic anxiety in humans with bilateral amygdala lesions: pharmacological induction via cardio-respiratory interoceptive pathways. *J Neurosci* 2016;36:3559-66.
61. Wiest G, Lehner-Baumgartner E, Baumgartner C. Panic attacks in an individual with bilateral selective lesions of the amygdala. *Arch Neurol* 2006;63:1798-801.
62. Adolphs R. What does the amygdala contribute to social cognition? *Ann N Y Acad Sci* 2010;1191:42-61.
63. Janak PH, Tye KM. From circuits to behaviour in the amygdala. *Nature* 2015;517:284-92.
64. Murray Sherman S, Guillery RW. Two types of thalamic relay. In: Murray Sherman S, Guillery RW, editors. *Exploring the Thalamus and Its Role in Cortical Function*, 2nd ed. Cambridge (MA): MIT Press; 2001. Available: www.sciencedirect.com/science/article/pii/B9780123054609500228 (accessed 2016 June 12).
65. Saalman YB, Kastner S. Cognitive and perceptual functions of the visual thalamus. *Neuron* 2011;71:209-23.
66. Gasquoine PG. Localization of function in anterior cingulate cortex: from psychosurgery to functional neuroimaging. *Neurosci Biobehav Rev* 2013;37:340-8.
67. Shackman AJ, Salomons TV, Slagter HA, et al. The integration of negative affect, pain, and cognitive control in the cingulate cortex. *Nat Rev Neurosci* 2011;12:154-67.
68. Shafer AT, Matveychuk D, Penney T, et al. Processing of emotional distraction is both automatic and modulated by attention: evidence from an event-related fMRI investigation. *J Cogn Neurosci* 2012;24:1233-52.
69. Straube T, Schmidt S, Weiss T, et al. Dynamic activation of the anterior cingulate cortex during anticipatory anxiety. *Neuroimage* 2009;44:975-81.
70. Etkin A, Büchel C, Gross JJ. The neural bases of emotion regulation. *Nat Rev Neurosci* 2015;16:693-700.
71. Lane RD, Chua PM-L, Dolan RJ. Common effects of emotional valence, arousal and attention on neural activation during visual processing of pictures. *Neuropsychologia* 1999;37:989-97.
72. Sabatinelli D, Fortune EE, Li Q, et al. Emotional perception: meta-analyses of face and natural scene processing. *Neuroimage* 2011;54:2524-33.
73. Vuilleumier P, Driver J. Modulation of visual processing by attention and emotion: windows on causal interactions between human brain regions. *Philos Trans R Soc Lond B Biol Sci* 2007;362:837-55.
74. Vuilleumier P. How brains beware: neural mechanisms of emotional attention. *Trends Cogn Sci* 2005;9:585-94.
75. Bandelow B. Panic and Agoraphobia Scale (PAS). Ashland (OH): Hogrefe & Huber Publishers; 1997.
76. Hautzinger M, Bailer M, Worall H, et al. Beck-Depressions-Inventar (BDI). Testhandbuch. 2. Auflage. Bern: Hans Huber; 1995.

Service Information

Subscriptions and sales

Annual subscriptions are available at the following rates in 2018 (Canadian customers pay in Canadian funds and add applicable taxes; US and rest of world pay in US funds): Can\$244 or US\$281 individuals, Can\$374 or US\$437 institutions, Can\$47 for Canadian students or residents, and Can\$36 or US\$47 for single or back issues. For more information or to subscribe, please contact the Joule Inc. Subscription Office, PO box 830350, Birmingham AL 35283-0350; phone 800 633-4931 (Canada and Continental US only) or 205 995-1567; fax 205 995-1588; cma@subscriptionoffice.com.

Change of address

Subscribers should send the new address and the effective date of change to the CMA Member Service Centre at cmamsc@cma.ca. Allow at least 8 weeks' notice to ensure uninterrupted service.

Reprints

Commercial and author reprints can be purchased through Sheridan Press. To purchase commercial article reprints and ePrints, or to request a quote, please contact Hope Robinson, Sheridan Content Sales Specialist; hope.robinson@sheridan.com. Authors can order reprints by submitting an author reprint order form available at the Sheridan Press Electronic

Order Centre at www.sheridan.com/cma/eoc or by contacting Lori Laughman, Customer Service Representative; lori.laughman@sheridan.com.

Permissions

Copyright for all material is held by Joule Inc., a wholly owned subsidiary of the Canadian Medical Association. We are a member of Access Copyright, The Canadian Copyright Licensing Agency, who may grant organizations and individuals, on our behalf, the right to respond to permissions requests. Please submit your request to Access Copyright using the online permission request service at <http://discovery.accesscopyright.ca>. For more information on licensing or to obtain a price estimate, visit www.accesscopyright.ca/permissions/

Microform, abstracting and indexing

The journal is abstracted or indexed in: MEDLINE/Index Medicus, Current Contents, Bioresearch Index, Social Sciences Citation Index, Biological Abstracts, EMBASE/Excerpta Medica, e-psyche, Mental Health Abstracts, ISI Web of Science, Child Development Abstracts and Bibliography, Standard Periodical Directory, Arts and Humanities Citation Index, Research Alert (formerly Automated Subject Citation Alert), Criminal Justice Abstracts, Neuroscience Citation Index, Psychological Abstracts, Sociological Abstracts. *JPN* is available on microform from ProQuest Information and Learning, PO Box 1346, Ann Arbor MI 48106-1346; il.proquest.com.