

# Neurobiological commonalities and distinctions among 3 major psychiatric disorders: a graph theoretical analysis of the structural connectome

Shuai Wang, MD; Gaolang Gong, PhD; Suyu Zhong, PhD; Jia Duan, MD; Zhiyang Yin, MD; Miao Chang, PhD; Shengnan Wei, MD, PhD; Xiaowei Jiang, MD; Yifang Zhou, MD; Yanqing Tang, MD, PhD; Fei Wang, MD, PhD

Published online on Aug. 1, 2019; subject to revision

**Background:** White matter network alterations have increasingly been implicated in major depressive disorder, bipolar disorder and schizophrenia. The aim of this study was to identify shared and distinct white matter network alterations among the 3 disorders. **Methods:** We used analysis of covariance, with age and gender as covariates, to investigate white matter network alterations in 123 patients with schizophrenia, 123 with bipolar disorder, 124 with major depressive disorder and 209 healthy controls. **Results:** We found significant group differences in global network efficiency ( $F = 3.386$ ,  $p = 0.018$ ), nodal efficiency ( $F = 8.015$ ,  $p < 0.001$  corrected for false discovery rate [FDR]) and nodal degree ( $F = 5.971$ ,  $p_{\text{FDR}} < 0.001$ ) in the left middle occipital gyrus, as well as nodal efficiency ( $F = 6.930$ ,  $p_{\text{FDR}} < 0.001$ ) and nodal degree ( $F = 5.884$ ,  $p_{\text{FDR}} < 0.001$ ) in the left postcentral gyrus. We found no significant alterations in patients with major depressive disorder. Post hoc analyses revealed that compared with healthy controls, patients in the schizophrenia and bipolar disorder groups showed decreased global network efficiency, nodal efficiency and nodal degree in the left middle occipital gyrus. Furthermore, patients in the schizophrenia group showed decreased nodal efficiency and nodal degree in the left postcentral gyrus compared with healthy controls. **Limitations:** Our findings could have been confounded in part by treatment differences. **Conclusion:** Our findings implicate graded white matter network alterations across the 3 disorders, enhancing our understanding of shared and distinct pathophysiological mechanisms across diagnoses and providing vital insights into neuroimaging-based methods for diagnosis and research.

## Introduction

Schizophrenia, bipolar disorder and major depressive disorder (MDD) are among the most common diseases in psychiatric nosology. There is increasing evidence for shared features among these disorders with respect to clinical characteristics<sup>1</sup> and genetic,<sup>2</sup> immune,<sup>3</sup> endocrine<sup>4</sup> and neural alterations.<sup>5,6</sup> Frequent difficulties in distinguishing these 3 disorders has resulted in obstacles to appropriate treatment.

Convergent evidence from postmortem, genetic and MRI studies underscores the significance of white matter pathology across the 3 disorders. Postmortem studies have shown white matter abnormalities in oligodendrocyte density and oligodendrocyte function, and in myelin abnormalities, across the 3 disorders,<sup>7,8</sup> and these findings are further supported by shared risk genes involved in oligodendrocyte

function and myelin.<sup>9-11</sup> Abnormal oligodendrocyte function and demyelination have been associated with alterations in white matter integrity in the 3 disorders.<sup>7,12,13</sup> Moreover, MRI studies have found commonalities in alterations in white matter integrity among schizophrenia, bipolar disorder and MDD,<sup>14-19</sup> with more prominent alterations in schizophrenia and bipolar disorder than in MDD.<sup>20</sup>

Increasing attention has been focused on white matter network alterations in schizophrenia, bipolar disorder and MDD. Many studies have used graph theory to describe white matter network architecture.<sup>21</sup> Using graph theory, network architecture can be assessed using different topological metrics, such as global efficiency, nodal efficiency and nodal degree. For example, Wang and colleagues<sup>22</sup> found reduced global efficiency of the white matter network in schizophrenia, and another investigation found that reduced nodal

**Correspondence to:** F. Wang, Department of Psychiatry and Radiology, First Affiliated Hospital, China Medical University, 155 Nanjing North Street, Heping District, Shenyang, Liaoning, 110001, PR China; fei.wang@cmu.edu.cn; Y. Tang, Department of Psychiatry and Geriatrics, First Affiliated Hospital, China Medical University, 155 Nanjing North Street, Heping District, Shenyang, Liaoning, 110001, PR China; yanqingtang@163.com

Submitted Sep. 12, 2018; Revised Jan. 25, 2019; Accepted Mar. 15, 2019

DOI: 10.1503/jpn.180162

efficiency in the hippocampus was linked to schizophrenia.<sup>23</sup> In bipolar disorder, reduced global efficiency and reduced nodal efficiency in the left inferior frontal gyrus have also been found.<sup>24,25</sup> However, network alterations are less clear in MDD.<sup>26–28</sup> A previous study found significant global and nodal topological alterations at an uncorrected threshold, but significance was lost with correction for multiple comparisons.<sup>26</sup> Neuroimaging studies have focused primarily on individual disorders, some comparing 2 disorders. Comparison studies involving schizophrenia, bipolar disorder and MDD are rare but can offer substantial insight into the development and pathophysiology of these psychiatric disorders.

In this study, we used graph theory analysis to examine white matter network architectures in schizophrenia, bipolar disorder, MDD and healthy controls, with particular interest in shared and distinct alterations among the 3 disorders. We hypothesized that shared white matter alterations would occur in a gradient manner, with schizophrenia > bipolar disorder > MDD in terms of extent of change.

## Methods

### *Participants and clinical assessment*

A total of 579 individuals (age 18–45 years) participated in this study, including 124 patients with MDD, 123 with bipolar disorder and 123 with schizophrenia, as well as 209 healthy controls. All patients with schizophrenia, bipolar disorder and MDD were recruited from the outpatient and inpatient Department of Psychiatry, First Affiliated Hospital of China Medical University, and Shenyang Mental Health Centre, Shenyang, China. Diagnosis was determined by 2 trained clinical psychiatrists according to the Structured Clinical Interview for DSM-IV Axis I Disorders (SCID). All patients met the diagnostic criteria of the DSM-IV for schizophrenia, bipolar disorder or MDD, without any other Axis I disorders. Healthy controls were recruited from the local region via advertisement. Each control participant was assessed using the non-patient version of the SCID to confirm that they had no current or lifetime Axis I disorder or history of Axis I disorders in first-degree relatives, as determined by a detailed family history.

Exclusion criteria for all participants included substance/alcohol abuse or dependence, or a concomitant major medical disorder; significant pathological change found via  $T_1$ - and  $T_2$ -weighted MRI; a history of head trauma with loss of consciousness for  $\geq 5$  minutes, or any neurologic disorder; any infectious disease, such as HIV/AIDS or severe acute respiratory syndrome; any MRI contraindications; or suboptimal imaging data quality.

All participants' symptoms were measured using the 17-item version of the Hamilton Depression Rating Scale, the Hamilton Anxiety Rating Scale, the Young Mania Rating Scale and the Brief Psychiatric Rating Scale.

This study was approved by the Medical Science Research Ethics Committee of First Affiliated Hospital of China Medical University. After receiving a detailed description of this study, every participant provided written informed consent.

### *MRI acquisition*

We performed MRI scans using a 3.0 T GE Sigma system (General Electric) with a standard 8-channel head coil at First Affiliated Hospital of China Medical University. We acquired  $T_1$  images using a 3-dimensional fast-spoiled gradient echo sequence with the following parameters: repetition time 7.1 ms, echo time 3.2 ms, image matrix  $240 \times 240$ , field of view  $240 \times 240$  mm<sup>2</sup>, 176 contiguous slices of 1 mm without gap, voxel size  $1.0 \times 1.0 \times 1.0$  mm<sup>3</sup>. We acquired diffusion tensor imaging (DTI) using a single short spin echo planar imaging sequence with the following parameters: repetition time 17000 ms, echo time 85.4 ms, image matrix  $120 \times 120$ , field of view  $240 \times 240$  mm<sup>2</sup>, 65 contiguous slices of 2 mm without gap, and 25 noncollinear directions ( $b = 1000$  s/mm<sup>2</sup>), together with an axial acquisition without diffusion weighting ( $b = 0$ ), voxel size  $2.0 \times 2.0 \times 2.0$  mm<sup>3</sup>. Two neuroradiologists with more than 3 years of experience interpreting neuroradiology images checked image quality.

### *MRI preprocessing*

The DTI data set was preprocessed using PANDA.<sup>29</sup> Briefly, data preprocessing included brain extraction, correction for eddy-current distortion and simple head motion, correction for b-matrix and computation for diffusion tensor and fractional anisotropy.

### *Network construction*

The construction of the white matter network was implemented in PANDA. The procedure consisted of 2 basic steps: definition of nodes and definition of edges.

#### **Network node definition**

We used an automated anatomic labelling (AAL) atlas to parcellate the brain's cortical and subcortical structures into 90 regions (pons and cerebellum excluded).<sup>30</sup> Each region represented a node of the white matter network. Briefly, the procedure included the following steps. First, the structural image (i.e.,  $T_1$ -weighted image) of each participant was coregistered to their  $b_0$  image in the DTI native space by linear transformation. Next, the coregistered image was nonlinearly normalized to the  $T_1$  template of ICBM152 in the Montreal Neurological Institute (MNI) space, resulting in a nonlinear transformation. Finally, the inverse transformation was applied to warp the AAL template from the MNI space to each participant's DTI native space.

#### **Network edge definition**

We implemented reconstruction of whole-brain white matter tracts using the deterministic fibre assignment continuous tracking algorithm.<sup>31</sup> If the turn angle of a fibre tract was greater than 45° or if any voxel of a fibre's fractional anisotropy became less than 0.2, the tracking procedure would be terminated. We determined that an edge existed if at least 3 tracts had terminal points between the 2 nodes. In this

study, we defined the weight of the edges by the mean fractional anisotropy value (an index for evaluating white matter fibre integrity<sup>32</sup>) of the white matter fibres between 2 nodes. Using these criteria, we obtained a  $90 \times 90$  fractional anisotropy weighted metric for each participant.

We used PANDA to check the quality of the registrations, such as the registration of DTI to  $T_1$ , the registration of  $T_1$  to MNI space and the registration of DTI to MNI space. We used TrackVis ([www.trackvis.org](http://www.trackvis.org)) to check the quality of deterministic fibre tracking.

### Network metrics

We computed topological metrics of the white matter network using graph theory. Because previous investigations suggested that degree and efficiency were associated with mental disorders,<sup>33–36</sup> we used GRETNA ([www.nitrc.org/projects/gretna/](http://www.nitrc.org/projects/gretna/)) to calculate them.<sup>37</sup> Brief descriptions and formulas are as follows.

#### Global network efficiency

In a complex network, the shortest path length is defined as the smallest sum of the edges throughout all possible paths from nodes  $i$  to  $j$  in the graph.<sup>38</sup> Global network efficiency is defined as the mean of the inverse of the shortest path length for all nodes in the network. For a given network  $G$  with  $N$  nodes, global network efficiency ( $E_{\text{glob}}^G$ ) is:

$$E_{\text{glob}}^G = \frac{1}{N(N-1)} \sum_{i \in G} \sum_{j \neq i \in G} \frac{1}{L_{ij}}$$

where  $L_{ij}$  is the shortest path length between node  $i$  and node  $j$ , and  $N$  is the number of nodes of the network  $G$ . Global network efficiency reflects the capacity to facilitate parallel information transfer within the entire network.

#### Local network efficiency

For a given node  $i$ ,  $G_i$  is the subgraph containing the neighbours of node  $i$  (the nodes directly connected with node  $i$ ) and the edges among them (Appendix 1, Figure S1, available at [jpn.ca/180162-a1](http://jpn.ca/180162-a1)). The local efficiency of node  $i$  ( $E_{\text{loc}}^i$ ) is defined as the global efficiency of  $G_i$ .

$$E_{\text{loc}}^i = E_{\text{glob}}^{G_i}$$

The local network efficiency ( $E_{\text{loc}}^G$ ) is described as the mean of the local efficiencies for all nodes in the network:<sup>38</sup>

$$E_{\text{loc}}^G = \frac{1}{N} \sum_{i \in G} E_{\text{loc}}^i$$

Local network efficiency corresponds to the ability to tolerate faults in the whole network.

#### Nodal degree

The nodal degree of a given node  $i$  is defined as the number of edges attached to the node. It indicates a node's direct connection with other nodes in the network.<sup>39</sup>

#### Nodal efficiency

The nodal efficiency of a given node  $i$  is defined as the average inverse of the shortest path length between node  $i$  and all other nodes in the network<sup>39</sup>

$$E_{\text{nodal}}^G(i) = \frac{1}{N-1} \sum_{i \neq j \in G} \frac{1}{L_{ij}}$$

where  $N$  represents the number of nodes for the whole network. Nodal efficiency represents the capacity of node  $i$  to communicate with other nodes in the network.

### Statistical analysis

To compare demographic characteristics and clinical symptoms, we performed analysis of variance and  $\chi^2$  tests. We implemented analysis of covariance (ANCOVA), with age and sex as covariates, to evaluate differences between white matter network topological metrics among the 4 groups. We performed post hoc pair-wise analyses to detect significant group effects in the ANCOVA. We performed partial correlation analyses, with age and sex as covariates, to investigate the relationships between the white matter network metrics and clinical symptoms. We set statistical significance for the analysis of demographic and clinical characteristics, as well as network metrics, at  $p < 0.05$ . For analyses involving nodal network metrics, we applied false discovery rate (FDR) correction for multiple comparisons (the number of tests was 90, and significance was set to a corrected  $p_{\text{FDR}} < 0.05$ ).

## Results

### Demographic and clinical characteristics

Detailed demographic and clinical data for the participants are listed in Table 1. We found no significant between-group differences in terms of age, sex or first-episode status. Illness duration ( $p = 0.002$ ) and medication status ( $p < 0.001$ ) were significantly different between the schizophrenia, bipolar disorder and MDD groups. We also found significant differences in scores among the 3 disorders on the Hamilton Depression Rating Scale, Hamilton Anxiety Rating Scale, Young Mania Rating Scale and Brief Psychiatric Rating Scale (all  $p < 0.001$ ).

### Global and local network efficiency among the 4 groups

We found a significant difference in global network efficiency among the 4 groups ( $p = 0.018$ ), and post hoc analysis revealed that the schizophrenia ( $p = 0.003$ ) and bipolar disorder ( $p = 0.047$ ) groups had significantly decreased global network efficiency compared with the healthy controls. We found no

significant difference in local network efficiency among the 4 groups ( $p = 0.095$ ; Table 2 and Fig. 1).

#### Nodal efficiency and degree among the 4 groups

The ANCOVA showed significant between-group effects in the nodal degree of the left middle occipital gyrus and the left postcentral gyrus (both  $p_{FDR} < 0.001$ ). Post hoc analysis revealed that compared with healthy controls, patients with

schizophrenia showed decreased nodal degree in the left middle occipital gyrus ( $p = 0.005$ ) and left postcentral gyrus ( $p < 0.001$ ), and patients with bipolar disorder showed decreased nodal degree in the left middle occipital gyrus ( $p < 0.001$ ).

Similarly, we found significant group differences in the nodal efficiency of the left middle occipital gyrus ( $p_{FDR} < 0.001$ ) and the left postcentral gyrus ( $p_{FDR} < 0.001$ ). Post hoc analysis revealed that compared with healthy controls, patients with schizophrenia showed decreased nodal efficiency

**Table 1: Demographic and clinical characteristics**

Characteristic	Group; mean $\pm$ SD or no. (%)				$F/\chi^2$	$p$ value	Post hoc significance
	Control ( $n = 209$ )	MDD ( $n = 124$ )	BD ( $n = 123$ )	Schizophrenia ( $n = 123$ )			
<b>Demographic</b>							
Age at scan, yr	28.91 $\pm$ 7.58	29.67 $\pm$ 8.04	27.50 $\pm$ 6.50	28.15 $\pm$ 7.91	1.96*	0.12*	—
Male sex	85 (40.7)	37 (29.8)	55 (44.7)	49 (39.8)	6.34*	0.096*	—
<b>Clinical</b>							
Duration, mo	—	24.88 $\pm$ 39.83	47.12 $\pm$ 55.69	33.51 $\pm$ 44.78	6.48†	0.002†	MDD vs. BD: $p = 0.002$
First episode	—	99 (79.8)	48 (39.0)	70 (56.9)	42.72†	0.08†	—
Medication	—	69 (55.6)	93 (75.6)	82 (66.7)	11.0†	0.004†	—
Antipsychotic	—	4 (3.2)	10 (8.1)	68 (55.3)	—	—	—
Mood stabilizer	—	0 (0)	7 (5.7)	1 (0.8)	—	—	—
Antidepressant	—	60 (48.4)	14 (11.4)	0 (0)	—	—	—
Combination	—	5 (4.0)	62 (50.4)	13 (10.6)	—	—	—
HAM-D, total score‡	—	18.85 $\pm$ 9.81	10.54 $\pm$ 9.33	7.16 $\pm$ 6.66	50.47†	< 0.001†	MDD vs. SZ: $p < 0.001$ MDD vs. BD: $p < 0.001$ BD vs. SZ: $p = 0.021$
HAM-A, total score§	—	15.87 $\pm$ 10.68	7.59 $\pm$ 8.08	6.00 $\pm$ 7.08	36.44†	< 0.001†	MDD vs. SZ: $p < 0.001$ MDD vs. BD: $p < 0.001$
YMRS, total score¶	—	1.51 $\pm$ 3.44	6.63 $\pm$ 9.49	1.84 $\pm$ 4.97	19.27†	< 0.001†	MDD vs. BD: $p < 0.001$ BD vs. SZ: $p < 0.001$
BPRS, total score**	—	25.9 $\pm$ 6.83	25.36 $\pm$ 9.65	31.42 $\pm$ 11.93	10.93†	< 0.001†	MDD vs. SZ: $p < 0.001$ BD vs. SZ: $p < 0.001$

BD = bipolar disorder; BPRS = Brief Psychiatric Rating Scale; HAM-A = Hamilton Anxiety Scale; HAM-D = Hamilton Depression Rating Scale; MDD = major depressive disorder; SD = standard deviation; SZ = schizophrenia; YMRS = Young Mania Rating Scale.

\*Examination among the healthy control, major depressive disorder, bipolar disorder and schizophrenia groups.

†Examination among the major depressive disorder, bipolar disorder and schizophrenia groups.

‡Major depressive disorder,  $n = 122$ ; bipolar disorder,  $n = 116$ ; schizophrenia  $n = 91$ .

§Major depressive disorder,  $n = 115$ ; bipolar disorder,  $n = 113$ ; schizophrenia  $n = 77$ .

¶Major depressive disorder,  $n = 110$ ; bipolar disorder,  $n = 115$ ; schizophrenia  $n = 69$ .

\*\*Major depressive disorder,  $n = 78$ ; bipolar disorder,  $n = 73$ ; schizophrenia  $n = 116$ .

**Table 2: Differences in topological metrics across groups**

Topological metric	Brain region	Group; mean $\pm$ SD				$F_3$	$p$ value	Post hoc significance
		Control	MDD	BD	Schizophrenia			
Global network efficiency	—	0.221 $\pm$ 0.011	0.220 $\pm$ 0.011	0.218 $\pm$ 0.012	0.217 $\pm$ 0.015	3.386	0.018*	HC vs. BD: $p = 0.047$ HC vs. SZ: $p = 0.003$
Local network efficiency	—	0.303 $\pm$ 0.013	0.301 $\pm$ 0.014	0.303 $\pm$ 0.014	0.299 $\pm$ 0.017	2.130	0.095	—
Nodal degree	Left middle occipital gyrus	6.425 $\pm$ 1.511	6.215 $\pm$ 1.250	5.828 $\pm$ 1.431	5.949 $\pm$ 1.165	5.971	< 0.001*	HC vs. BD: $p < 0.001$ HC vs. SZ: $p = 0.005$
	Left postcentral gyrus	5.523 $\pm$ 1.242	5.445 $\pm$ 1.212	5.448 $\pm$ 1.126	4.970 $\pm$ 1.248	5.884	< 0.001*	HC vs. SZ: $p < 0.001$
Nodal efficiency	Left middle occipital gyrus	0.235 $\pm$ 0.021	0.231 $\pm$ 0.020	0.225 $\pm$ 0.023	0.226 $\pm$ 0.019	8.015	< 0.001*	HC vs. BD: $p < 0.001$ HC vs. SZ: $p < 0.001$
	Left postcentral gyrus	0.230 $\pm$ 0.020	0.229 $\pm$ 0.022	0.229 $\pm$ 0.019	0.220 $\pm$ 0.021	6.930	< 0.001*	HC vs. SZ: $p < 0.001$

BD = bipolar disorder; HC = healthy controls; SD, standard deviation; SZ = schizophrenia.

\*Significant at  $p < 0.05$ .

in the left middle occipital gyrus ( $p < 0.001$ ) and left postcentral gyrus ( $p < 0.001$ ), and patients with bipolar disorder showed decreased nodal efficiency in the left middle occipital gyrus ( $p < 0.001$ ; Table 2 and Fig. 2).

#### Correlations between topological metrics and clinical characteristics


Following FDR correction, we found no significant correlations between topological metrics and clinical characteristics.

### Discussion


To our knowledge, this is the first study to examine white matter network alterations across schizophrenia, bipolar disorder and MDD with a large sample. We found shared decreased global network efficiency, nodal efficiency and nodal degree in the left middle occipital gyrus in patients with schizophrenia and bipolar disorder. Specific to schizophrenia only, we observed significant decreased nodal efficiency and nodal degree in the left postcentral gyrus. Shared alterations were more prominent in schizophrenia and bipolar disorder, but greatest in schizophrenia. Post hoc analyses did not find significant changes in MDD. Our findings supported the idea that network alterations occur in a graded manner (schizophrenia > bipolar disorder > MDD).

Consistent with previous findings, this study found significant network alterations reflected in decreased global network efficiency in schizophrenia and bipolar disorder but not in MDD. Many graph theory studies have indicated that patients with schizophrenia show disrupted global white matter net-

work integration, including elevated clustering and reduced global network efficiency.<sup>33,35,36</sup> Studies have also found reduced white matter network integration in bipolar disorder.<sup>24,40,41</sup> For example, Collin and colleagues<sup>24</sup> compared patients with bipolar I disorder to healthy controls and found that patients exhibited reduced global network efficiency, which was associated with interhemispheric connectivity disruption. In contrast, findings in MDD have been mixed.<sup>26–28</sup> For example, Korgaonkar and colleagues<sup>26</sup> examined the brain network in MDD compared with healthy controls using 2 methods: network-based statistics and graph analysis. In their network-based statistics analyses, they found abnormalities mainly in the default mode network and in regions important for emotional and cognitive processing (including the thalamus, caudate and frontal cortex) in MDD. However, using graph analysis, they observed no significant network alterations in MDD compared with healthy controls. Nevertheless, the Korgaonkar graph theory findings were consistent with our own and could indicate fewer global alterations in MDD. The divergence of findings in MDD might originate from methodological differences and requires further investigation. Taken together, our findings were consistent with the literature and suggest similarities in disrupted global white matter network integration in both schizophrenia and bipolar disorder. The present study implicated nodal topological differences in the left hemisphere (i.e., the left middle occipital gyrus and left postcentral gyrus). Structural and functional lateralization of the human brain is well described in the literature:<sup>42–44</sup> the left hemisphere is dominant for the processing of language, and the right hemisphere is specialized for some nonverbal functions, such as spatial attention and face



**Fig. 1:** Each dot represents 1 participant's global network efficiency. \*Significant at  $p < 0.05$ . \*\*\*Significant at  $p < 0.005$ . BD = bipolar disorder; HC = healthy controls; MDD = major depressive disorder; SZ = schizophrenia.



**Fig. 2:** (A, C) Three-dimensional representations of the nodal degree and nodal efficiency between the 4 groups. The node size represents the mean nodal degree/nodal efficiency of each group. (B, D) Violin plots of the nodal degree and nodal efficiency between the 4 groups. Each dot represents 1 participant's nodal degree/nodal efficiency. \*Significant at  $p < 0.05$ . \*\*Significant at  $p < 0.01$ . \*\*\*Significant at  $p < 0.005$ . \*\*\*\*Significant at  $p < 0.001$ . BD = bipolar disorder; HC = healthy controls; MDD = major depressive disorder; MOG.L = left middle occipital gyrus; PoCG.L = left postcentral gyrus; SZ = schizophrenia.

processing. Previous studies have shown decreased language lateralization in psychiatric disorders.<sup>45–47</sup> In addition, language lateralization has been associated with clinical symptom severity.<sup>46</sup> Altered hemispheric lateralization could serve as a neural marker of psychiatric disorders.<sup>23,48</sup> Further, our findings may implicate the importance of the left hemisphere in schizophrenia and bipolar disorder.

In this study, we found reduced nodal efficiency and nodal degree in the middle occipital gyrus in patients with schizophrenia and bipolar disorder. The middle occipital gyrus is associated with visual information processing. Previous studies have indicated ubiquitous visual system dysfunction in schizophrenia and bipolar disorder.<sup>49,50</sup> Furthermore, Knöchel and colleagues<sup>51</sup> found that visual alterations were correlated with altered white matter integrity both in schizophrenia and bipolar disorder patients. In line with these findings, our results suggest that the reduced nodal degree and nodal efficiency of the middle occipital gyrus might be a common feature between schizophrenia and bipolar disorder.

In the present study, we observed decreased postcentral gyrus nodal degree and nodal efficiency only in patients with schizophrenia. The postcentral gyrus is known as the primary somatosensory cortex and plays a role in the representation of somatotopic information and the sensory reception of touch and kinesthesia.<sup>52</sup> Previous studies have found that schizophrenia patients have reduced white matter integrity of 2 major ascending tracts to the primary somatosensory cortex,<sup>53</sup> and reduced grey matter volume of the postcentral gyrus.<sup>54</sup> These findings are supported by decreased functional activation of the postcentral gyrus in schizophrenia.<sup>55,56</sup> Decreased nodal efficiency in the postcentral gyrus may be a distinct feature of schizophrenia, separating it from bipolar disorder and MDD.

### Limitations

Several limitations should be taken into account in this study. First, approximately 60% of the participants were taking anti-psychotic medications or antidepressants. To avoid possible drug confounds, future studies should recruit medication-naïve participants for the effects of psychotropic medication. Second, the AAL we used is the most frequently used atlas, but others are available and may be more refined in localizing findings.<sup>28</sup> Also, probabilistic tractography may be better than deterministic tractography in accounting for uncertainty in the data.<sup>57,58</sup> Further studies should aim to use a high-resolution atlas and probabilistic tractography to verify the results of this study.

### Conclusion

Our study found significant network alterations in schizophrenia and bipolar disorder but not in MDD. Patients with schizophrenia and bipolar disorder showed reduced global efficiency and nodal efficiency of the middle occipital gyrus. We observed alterations in the postcentral gyrus only in schizophrenia. Our findings were consistent with previous studies and supported more similarities between schizophrenia and bipolar disorder than with MDD. They also implicated more prominent network alterations in schizophrenia and bipolar

disorder than in MDD and supported a gradient of network alterations (schizophrenia > bipolar disorder > MDD). These findings inform future approaches to understanding the brain network across schizophrenia, bipolar disorder and MDD.

**Acknowledgements:** This work was supported by grants from the National Natural Science Foundation of China (81701336 to S. Wei, 81571311, 81071099 and 81271499 to Yanqing Tang, 81725005, 81571331 to Fei Wang), the National Key Research and Development Program (2016YFC1306900 to Yanqing Tang), the Liaoning Education Foundation (L2015591 to Shengnan Wei), the Liaoning Education Foundation (Pandeng Scholar, Fei Wang), the National Key Research and Development Program (2016YFC0904300 to Fei Wang), and the National High Tech Development Plan (863; 2015AA020513 to Fei Wang). The authors thank Dr. Fay Y. Womer, Department of Psychiatry, Washington University School of Medicine, and Dr. E. Kale Edmiston, Department of Psychiatry, University of Pittsburgh Medical Center, for their editing of this manuscript.

**Affiliations:** From the Department of Psychiatry, First Affiliated Hospital of China Medical University, Shenyang, Liaoning, China (S. Wang, Duan, Yin, Tang, F. Wang); the State Key Laboratory of Cognitive Neuroscience and Learning & IDG/McGovern Institute for Brain Research, Beijing Normal University, Beijing, China (Gong, Zhong); the Department of Radiology, First Affiliated Hospital of China Medical University, Shenyang, Liaoning, China (Chang, Wei, Jiang, F. Wang); the Brain Function Research Section, First Affiliated Hospital of China Medical University, Shenyang, Liaoning, China (S. Wang, Duan, Chang, Wei, Jiang, Zhou, Tang, F. Wang); and the Department of Gerontology, First Affiliated Hospital of China Medical University, Shenyang, Liaoning, China (Zhou, Tang).

**Competing interests:** None declared.

**Contributors:** S. Wang, G. Gong, Y. Tang and F. Wang designed the study. S. Wang, J. Duan, Z. Yin, X. Jiang and Y. Zhou acquired the data, which S. Wang, G. Gong, S. Zhong, J. Duan, Z. Yin, M. Chang and S. Wei analyzed. S. Wang, Y. Tang and F. Wang wrote the article, which all authors reviewed. All authors approved the final version to be published and can certify that no other individuals not listed as authors have made substantial contributions to the paper.

### References

1. Barch DM, Sheffield JM. Cognitive impairments in psychotic disorders: common mechanisms and measurement. *World Psychiatry* 2014;13:224–32.
2. Consortium C-DGotPG. Identification of risk loci with shared effects on five major psychiatric disorders: a genome-wide analysis. *Lancet* 2013;381:1371–9.
3. Reus GZ, Fries GR, Stertz L, et al. The role of inflammation and microglial activation in the pathophysiology of psychiatric disorders. *Neuroscience* 2015;300:141–54.
4. Garcia-Rizo C, Kirkpatrick B, Fernandez-Egea E, et al. Abnormal glycemic homeostasis at the onset of serious mental illnesses: a common pathway. *Psychoneuroendocrinology* 2016;67:70–5.
5. Network PASoPGC. Psychiatric genome-wide association study analyses implicate neuronal, immune and histone pathways. *Nat Neurosci* 2015;18:199–209.
6. Goodkind M, Eickhoff SB, Oathes DJ, et al. Identification of a common neurobiological substrate for mental illness. *JAMA Psychiatry* 2015;72:305–15.
7. Tham MW, Woon PS, Sum MY, et al. White matter abnormalities in major depression: evidence from post-mortem, neuroimaging and genetic studies. *J Affect Disord* 2011;132:26–36.
8. Mighdoll MI, Tao R, Kleinman JE, et al. Myelin, myelin-related disorders, and psychosis. *Schizophr Res* 2015;161:85–93.
9. Perlstein MD, Chohan MR, Coman IL, et al. White matter abnormalities in 22q11.2 deletion syndrome: preliminary associations with the Nogo-66 receptor gene and symptoms of psychosis. *Schizophr Res* 2014;152:117–23.

10. Whalley HC, Dimitrova R, Sprooten E, et al. Effects of a balanced translocation between chromosomes 1 and 11 disrupting the droshoph Inf ServC1 Locus on white matter integrity. *PLoS One* 2015; 10: e0130900
11. Mallas EJ, Carletti F, Chaddock CA, et al. Genome-wide discovered psychosis-risk gene ZNF804A impacts on white matter microstructure in health, schizophrenia and bipolar disorder. *Peer J* 2016; 4:e1570.
12. Mahon K, Burdick KE, Szeszko PR. A role for white matter abnormalities in the pathophysiology of bipolar disorder. *Neurosci Biobehav Rev* 2010;34:533-54.
13. Herring NR, Konradi C. Myelin, copper, and the cuprizone model of schizophrenia. *Front Biosci* 2011;3:23-40.
14. Jovicich J, Marizzoni M, Sala-Llonch R, et al. Brain morphometry reproducibility in multi-center 3 T MRI studies: a comparison of cross-sectional and longitudinal segmentations. *Neuroimage* 2013; 83:472-84.
15. Skudlarski P, Schretlen DJ, Thaker GK, et al. Diffusion tensor imaging white matter endophenotypes in patients with schizophrenia or psychotic bipolar disorder and their relatives. *Am J Psychiatry* 2013;170:886-98.
16. Mamah D, Wen J, Luo J, et al. Subcomponents of brain T2\* relaxation in schizophrenia, bipolar disorder and siblings: a Gradient Echo Plural Contrast Imaging (GEPCI) study. *Schizophr Res* 2015; 169:36-45.
17. Nazeri A, Chakravarty MM, Felsky D, et al. Alterations of superficial white matter in schizophrenia and relationship to cognitive performance. *Neuropsychopharmacology* 2013;38:1954-62.
18. Mahon K, Wu J, Malhotra AK, et al. A voxel-based diffusion tensor imaging study of white matter in bipolar disorder. *Neuropsychopharmacology* 2009;34:1590-600.
19. Chang M, Womer FY, Edmiston EK, et al. Neurobiological commonalities and distinctions among three major psychiatric diagnostic categories: a structural MRI study. *Schizophr Bull* 2018; 44:65-74.
20. Kempton MJ, Salvador Z, Munafo MR, et al. Structural neuroimaging studies in major depressive disorder. Meta-analysis and comparison with bipolar disorder. *Arch Gen Psychiatry* 2011;68:675-90.
21. Bullmore E, Sporns O. Complex brain networks: graph theoretical analysis of structural and functional systems. *Nat Rev Neurosci* 2009;10:186-98.
22. Wang Q, Su TP, Zhou Y, et al. Anatomical insights into disrupted small-world networks in schizophrenia. *Neuroimage* 2012; 59:1085-93.
23. Sun Y, Chen Y, Collinson SL, et al. Reduced hemispheric asymmetry of brain anatomical networks is linked to schizophrenia: a connectome study. *Cereb Cortex* 2017;27:602-15.
24. Collin G, van den Heuvel MP, Abramovic L, et al. Brain network analysis reveals affected connectome structure in bipolar I disorder. *Hum Brain Mapp* 2016;37:122-34.
25. Liu H, Zhao K, Shi J, et al. Topological properties of brain structural networks represent early predictive characteristics for the occurrence of bipolar disorder in patients with major depressive disorder: a 7-year prospective longitudinal study. *Front Psychiatry* 2018;9:704.
26. Korgaonkar MS, Fornito A, Williams LM, et al. Abnormal structural networks characterize major depressive disorder: a connectome analysis. *Biol Psychiatry* 2014;76:567-74.
27. Qin J, Wei M, Liu H, et al. Abnormal brain anatomical topological organization of the cognitive-emotional and the frontoparietal circuitry in major depressive disorder. *Magn Reson Med* 2014;72:1397-407.
28. Ye M, Qing P, Zhang K, et al. Altered network efficiency in major depressive disorder. *BMC Psychiatry* 2016;16:450.
29. Cui Z, Zhong S, Xu P, et al. PANDA: a pipeline toolbox for analyzing brain diffusion images. *Front Hum Neurosci* 2013;7:42.
30. Tzourio-Mazoyer N, Landeau B, Papathanassiou D, et al. Automated anatomical labeling of activations in SPM using a macroscopic anatomical parcellation of the MNI MRI single-subject brain. *Neuroimage* 2002;15:273-89.
31. Mori S, Crain BJ, Chacko VP, et al. Three-dimensional tracking of axonal projections in the brain by magnetic resonance imaging. *Ann Neurol* 1999;45:265-9.
32. Beaulieu C. The basis of anisotropic water diffusion in the nervous system: a technical review. *NMR Biomed* 2002;15:435-55.
33. Collin G, de Nijs J, Hulshoff Pol HE, et al. Connectome organization is related to longitudinal changes in general functioning, symptoms and IQ in chronic schizophrenia. *Schizophr Res* 2016; 173:166-73.
34. van den Heuvel MP, Sporns O. Rich-club organization of the human connectome. *J Neurosci* 2011;31:15775-86.
35. van den Heuvel MP, Sporns O, Collin G, et al. Abnormal rich club organization and functional brain dynamics in schizophrenia. *JAMA Psychiatry* 2013;70:783-92.
36. Zalesky A, Fornito A, Seal ML, et al. Disrupted axonal fiber connectivity in schizophrenia. *Biol Psychiatry* 2011;69:80-9.
37. Wang JH, Wang XD, Xia MR, et al. GRETNA: a graph theoretical network analysis toolbox for imaging connectomics. *Front Hum Neurosci* 2015;9:386.
38. Latora V, Marchiori M. Efficient behavior of small-world networks. *Phys Rev Lett* 2001;87.
39. Achard S, Bullmore ET. Efficiency and cost of economical brain functional networks. *PLoS Comput Biol* 2007;3:174-83.
40. O'Donoghue S, Holleran L, Cannon DM, et al. Anatomical dysconnectivity in bipolar disorder compared with schizophrenia: a selective review of structural network analyses using diffusion MRI. *J Affect Disord* 2017;209:217-28.
41. Leow A, Ajilore O, Zhan L, et al. Impaired inter-hemispheric integration in bipolar disorder revealed with brain network analyses. *Biol Psychiatry* 2013;73:183-93.
42. Galaburda AM, LeMay M, Kemper TL, et al. Right-left asymmetries in the brain. *Science* 1978;199:852-6.
43. Toga AW, Thompson PM. Mapping brain asymmetry. *Nat Rev Neurosci* 2003;4:37-48.
44. Herve PY, Zago L, Petit L, et al. Revisiting human hemispheric specialization with neuroimaging. *Trends Cogn Sci* 2013;17:69-80.
45. Razafimandimby A, Maiza O, Herve PY, et al. Stability of functional language lateralization over time in schizophrenia patients. *Schizophr Res* 2007;94:197-206.
46. Sommer IE, Ramsey NF, Kahn RS. Language lateralization in schizophrenia, an fMRI study. *Schizophr Res* 2001;52:57-67.
47. Bleich-Cohen M, Hendler T, Kotler M, et al. Reduced language lateralization in first-episode schizophrenia: an fMRI index of functional asymmetry. *Psychiatry Res* 2009;171:82-93.
48. Yang C, Zhong S, Zhou X, et al. The abnormality of topological asymmetry between hemispheric brain white matter networks in Alzheimer's disease and mild cognitive impairment. *Front Aging Neurosci* 2017;9:261.
49. Xu GY, Lin KG, Rao DP, et al. Neuropsychological performance in bipolar I, bipolar II and unipolar depression patients: a longitudinal, naturalistic study. *J Affect Disord* 2012;136:328-39.
50. Zheng W, Tang LR, Correll CU, et al. Frequency and correlates of distant visual impairment in patients with schizophrenia, bipolar disorder, and major depressive disorder. *East Asian Arch Psychiatry* 2015;25:115-21.
51. Knochel C, Stablein M, Storchak H, et al. Multimodal assessments of the hippocampal formation in schizophrenia and bipolar disorder: evidences from neurobehavioral measures and functional and structural MRI. *Neuroimage Clin* 2014;6:134-44.
52. Ferro A, Roiz-Santianez R, Ortiz-Garcia de la Foz V, et al. A cross-sectional and longitudinal structural magnetic resonance imaging study of the post-central gyrus in first-episode schizophrenia patients. *Psychiatry Res* 2015;231:42-9.
53. Douaud G, Smith S, Jenkinson M, et al. Anatomically related grey and white matter abnormalities in adolescent-onset schizophrenia. *Brain* 2007;130:2375-86.
54. Job DE, Whalley HC, McConnell S, et al. Structural gray matter differences between first-episode schizophrenics and normal controls using voxel-based morphometry. *Neuroimage* 2002;17:880-9.
55. Wei Y, Chang M, Womer FY, et al. Local functional connectivity alterations in schizophrenia, bipolar disorder, and major depressive disorder. *J Affect Disord* 2018;236:266-73.
56. Zhuo C, Zhu J, Wang C, et al. Brain structural and functional dissociated patterns in schizophrenia. *BMC Psychiatry* 2017;17:45.
57. Descoteaux M, Deriche R, Knosche TR, et al. Deterministic and probabilistic tractography based on complex fibre orientation distributions. *IEEE Trans Med Imaging* 2009;28:269-86.
58. Jones DK. Tractography gone wild: probabilistic fibre tracking using the wild bootstrap with diffusion tensor MRI. *IEEE Trans Med Imaging* 2008;27:1268-74.