

**Appendix 1** to Zhao W, Zimmerman K, Zhou X, et al. Impaired cognitive performance under psychosocial stress in cannabis-dependent males is associated with attenuated precuneus activity. *J Psychiatry Neurosci* 2019.

DOI: 10.1503/jpn.190039

© Joule Inc.

*Online appendices are unedited and posted as supplied by the authors.*

## **Impaired cognitive performance under psychosocial stress in cannabis-dependent males is associated with attenuated precuneus activity**

Zhao et al.,

Correspondence: ben\_becker@gmx.de

### **Supplemental Materials**

#### **Data quality assessment and exclusion of participants**

Two cannabis users were excluded due to excessive use of other illicit substances. After initial data quality assessments, data from four cannabis users and three controls were excluded due to excessive head movement (> 3.5 mm) during fMRI. Two controls were excluded because they consistently rated subjective stress during the control as well as stress condition very low (stress experience was assessed using a 1-8 scale; one subject consistently rated the stress experience as 1 the other consistently as 2, indicating that the paradigm did not induce stress in these subjects). The final dataset thus included n = 28 dependent cannabis users and n = 23 healthy controls. Details are additionally visualized in the **Figure S1**.

#### **MRI data acquisition**

Data were acquired on a Siemens Trio 3T MRI system (Siemens, Erlangen, Germany). Functional data was acquired using a T2\* echo-planar imaging (EPI) BOLD sequence [repetition time (TR) = 2500 ms, echo time (TE) = 30 ms, 37 slices, slice thickness = 3.0 mm, no gap, voxel size = 2 × 2 × 3 mm, flip angle = 90°, field of view = 192 mm]. To exclude subjects with apparent brain pathologies and facilitate normalization high-resolution T1-weighted structural images were acquired (TR = 1660 ms, TE = 2540 ms, 208 slices, field of view = 256 mm, voxel size = 0.8 × 0.8 × 0.8 mm).

#### **fMRI data processing**

fMRI data were analyzed using SPM12 (Wellcome Department, London, UK, <http://www.fil.ion.ucl.ac.uk/spm/software/spm12>). The first five volumes were discarded to achieve magnet-steady images. Next, slice time correction was employed to correct for slice acquisition time and images were realigned using a six-parameter rigid body algorithm to correct for head movement. Subsequently, images were normalized using a two-step procedure that included co-registration with the T1 image, segmentation of the T1 image and application of the resulting transformation matrix to the functional time-series. Images were written out at 3 × 3 × 3 mm resolution and smoothed with a Gaussian kernel (FWHM, 8 mm).

**Appendix 1** to Zhao W, Zimmerman K, Zhou X, et al. Impaired cognitive performance under psychosocial stress in cannabis-dependent males is associated with attenuated precuneus activity. *J Psychiatry Neurosci* 2019.

DOI: 10.1503/jpn.190039

© Joule Inc.

*Online appendices are unedited and posted as supplied by the authors.*

### **fMRI BOLD level Results**

In line with previous studies, the paradigm induced widespread activity in psycho-social stress networks encompassing middle frontal regions, precuneus and posterior cingulate cortex (family-wise error correction  $p < 0.05$ ) (see also Dedovic et al., 2009; Wang et al., 2005). See also **Figure S2** showing the overlap between the stress network in the present study and the ROI mask generated from the Eckstein et al., 2014 sample.

### **Supplemental references**

Dedovic K, Rexroth M, Wolff E, Duchesne A, Scherling C, Beaudry T, Lue SD, Lord C, Engert V, Pruessner JC (2009). Neural correlates of processing stressful information: an event-related fMRI study. *Brain research* 1293, 49–60.

Eckstein M, Scheele D, Weber K, Stoffel-Wagner B, Maier W, Hurlmann R (2014). Oxytocin facilitates the sensation of social stress. *Human brain mapping* 35, 4741–4750.

Wang J, Rao H, Wetmore GS, Furlan PM, Korczykowski M, Dinges DF, Detre JA (2005). Perfusion functional MRI reveals cerebral blood flow pattern under psychological stress. *Proceedings of the National Academy of Sciences* 102, 17804–17809.


**Appendix 1** to Zhao W, Zimmerman K, Zhou X, et al. Impaired cognitive performance under psychosocial stress in cannabis-dependent males is associated with attenuated precuneus activity. *J Psychiatry Neurosci* 2019.

DOI: 10.1503/jpn.190039

© Joule Inc.

*Online appendices are unedited and posted as supplied by the authors.*

**Figure S1** Flow Charts for participants.



**Appendix 1** to Zhao W, Zimmerman K, Zhou X, et al. Impaired cognitive performance under psychosocial stress in cannabis-dependent males is associated with attenuated precuneus activity. *J Psychiatry Neurosci* 2019.

DOI: 10.1503/jpn.190039

© Joule Inc.

*Online appendices are unedited and posted as supplied by the authors.*

**Figure S2** Stress-related network in the current sample.

